

柒、參考文獻

- 呂鋒洲，林仁混。1991。基礎酵素學。聯經出版社，台北。
- 張文重。1976。蛋白質分解酵素(構造 功能 進化及應用)。國立編譯館，台北。
- 陳國誠。1989。微生物酵素工程學。藝軒出版社，台北。
- 賴威安。2000。*Bacillus* sp. P-6 中蛋白酶的生產與性質分析。國立中興大學食品科學系碩士論文。
- 張資奇。1999。蛋雞糞堆肥中篩選羽毛分解菌之研究。私立東海大學畜產學系研究所碩士論文。
- 李春蓮。2000。*Bacillus licheniformis* 聚木糖分解酵素之純化與生化特性探討。國立屏東科技大學食品科學系碩士論文。
- 賴滋漢，李秀。1976。食品分析與檢驗。精華出版社。
- 謝魁鵬，魏耀揮。1985。最新生物化學實驗。藝軒出版社，台北。
- 蘇遠志。1999。應用微生物學。華香園出版社，台北。
- 台灣統計年報。2002。
- Anwar, A. and M Saleemuddin. 1998. Alkaline proteases: a review. *Bioresource Technol.* 64:175-183.
- Barrett, A. J. and N. D. Rawlings. 1995. Families and clans of serine peptidases. *Arch. Environ. Microbiol.* 318(2):247-250.
- Beynon, R. J. and J. S. Bond. 1989. Proteolytic enzyme. Pp. 244-246. Oxford university Press Inc., New York.
- Bockle, B. and R. Muller. 1997. Reduction of disulfide bonds by *Streptomyces pactum* during growth on chicken feathers. *Appl. Environ. Microbiol.* 63(2):790-792.

- Bockle, B., B. Galunsky and R. Muller. 1995. Characterization of a keratinolytic serine proteinase from *Streptomyces pactum* DSM 40530. *Appl. Environ. Microbiol.* 61(10):3705-3710.
- Dalev, P. G and L. S. Simeonova. 1992. An enzyme biotechnology for the total utilization of leather wastes. *Biotech. Letters.* 14(6):531-534.
- Evans, K. L., J. Crowder and E. S. Miller. 2000. Subtilisins of *Bacillus spp.* hydrolyze keratin and allow growth on feathers. *Can. J. Microbiol.* 46:1004-1011.
- Ferrero, M. A., G. R. Castro, C. M. Abate and M. D. Baigori. 1996. Thermostable alkaline proteases of *Bacillus licheniformis* MIR29: isolation, production and characterization. *Microbiol Biotech.* 45: 327-332.
- Figueras, M. J., J. Guarro and L. Zaror. 1997. Ultrastructural aspects of hair digestion in black piedra infection. *J. Med. Vet. Mycol.* 35:1-6.
- Friedrich, A. B. and G. Antranikian. 1996. Keratin degradation by *Fervidobacterium pennavorans*, a novel thermophilic anaerobic species of the order *Thermotogales*. *Appl. Environ. Microbiol.* 62(8):2875-2882.
- Fujiwara, N. and A. Masui. 1993. Purification and properties of the highly thermostable alkaline protease from an alkalophilic and thermophilic *Bacillus* sp. *J. Biotech.* 30:245-256.
- Gupta, R., Q. K. Beg and P. Lorenz. 2002. Bacterial alkaline proteases: molecular approaches and industrial applications. *Appl. Microbiol. Biotechnol.* 59:15-32.
- Hazes, P. and B. W. Dijkstra. 1988. Model building of disulfide bonds in proteins with known three-dimensional structure. *Prot. Eng.* 2(2):119-125.
- Hoondal, G. S., R. P. Tiwari, N. Dahiya and Q. K. Beg. 2002. Microbial alkaline pectinases and their industrial applications: a review. *Appl. Microbiol Biotech.* 59: 409-418.

Kluskens, L. D., W. G. B. Voorhorst, R. J. Siezen, R. M. Schwerdtfeger, G. A. ntranikian, J. Oost and W. M. Vos. 2002. Molecular characterization of fervidolysin, a subtilisin-like serine protease from the thermophilic bacterium *Fervidobacterium pennivorans*. Extremophiles. 6: 185-194.

Kim, W., K Choi, Y Kim, H Park, J Choi, Y Lee, H Oh, I Kwon, and S Lee. 1996. Purification and characterization of a fibrinolytic enzyme produced from *Bacillus* sp. strain CK 11-4 screened from Chungkook-Jang. Appl. Environ. Microbiol. 62(7): 2482-2488.

Kobayashi, T., Y. Hakamada, S. Adachi, J. Hitomi, T. Yoshimatsu, K. Koike, S. kawai and S. Ito. 1996. Purification and properties of an alkaline protease from alkalophilic *Bacillus* sp. KSM-K16. Appl. Microbiol Biotech. 43:473-481.

Kumar, C. G. and H. Takagi. 1999. Microbial alkaline proteases: from a bioindustrial viewpoint. Biotech. Adv. 17:561-594.

Lehninger, A. L., D. L. Nelson and M. M. Cox. 1993. The three-dimensional structure of proteins. In: Principles of Biochemistry. 2nd ed. Worth Publishers, New York.

Lin, X., C. G. Lee, E. S. Casale and J. C. H. Shih. 1992. Purification and characterization of a keratinase from a feather-degrading *Bacillus licheniformis* strain. Appl. Environ. Microbiol. 58(10): 3271-3275.

Lin, X., D. W. Kelemen, E. S. Miller and J. C. H. Shih. 1995. Necleotide sequence and expression of kerA, the gene encoding a keratinolytic protease of *Bacillus licheniformis* PWD-1. Appl. Environ. Microbiol. 61(4):1469-1474.

Lin, X., J. C. H. Shih, and H. E. Swaisgood. 1996. Hydrolysis of feather keratin by immobilized keratinase. Appl. Environ. Microbiol. 62(11):4273-4275.

Lin, X., S. L. Lee, ES. Miller and J. C. H. Shh. 1997. Expression of *Bacillus licheniformis* PWD-1 keratinase gene in B. subtilis. J. Indust. Microbio & Biotech. 19: 134-138.

- Lin, X., S. W. Lee, H. D. Bae, J. A. Shelford and K. J. Cheng. 2001. Comparison of two feather-degrading *Bacillus licheniformis* strains. Asian-Aust. J. Anim. Sci. 14(12):1769-1774.
- Malviya, H. K., R. C. Rajak and S. K. Hasija. 1992. Synthesis and regulation of extracellular keratinase in three fungi isolated from the grounds of a gelatin factory, Jabalpur, India. Mycopathologia. 120:1-4.
- Nakanishi, T. and T. Yamamoto. 1974. Action and specificity of a *Streptomyces* alkalophilic proteinase. Agr. Biol. Chem. 38(12):2391-2397.
- Nam. G., D. W. Lee, H. S. Lee, N. J. Lee, B. C. Kim, E. A. Choe, J. K. Hwang, M. T. Suhartono, Y. R. Pyun. 2002. Native-feather degradation by *Fervidobacterium islandicum* AW-1, a newly isolated keratinase-producing thermophilic anaerobe. Arch. Microbiol. 178: 538-547.
- Onifade, A. A., N. A. Al-Sane, A. A. Al-Musallam and S. Al-Zarban. 1998. Potentials for biotechnological applications of keratin-degrading microorganisms and their enzymes for nutritional improvement of feathers and other keratins as livestock feed resources. Bioresource Technol. 66: 1-11.
- Rosen, H. 1957. A modified colorimetric analysis for amino acids. Arch. Biochem. Biophys. 67:10-15.
- Rozs, M., L. Manczinger, Cs. Vagvolgyi, F. Kevei. 2001. Secretion of a trypsin-like thiol protease by a new keratinolytic strain of *Bacillus licheniformis*. FEMS Microbiol. Lett. 205:221-224.
- Shih, J. C. H. 1993. Recent development in poultry waste digestion and feather utilization—a review. Poult. Sci. 72:1617-1620.
- Siezen R. J. and J. A. M. Leunissen. 1997. Subtilases: the superfamily of subtilisin-like serine proteases. Prot. Sci. 6: 501-523.
- Takagi, H., T. Takahashi, H. Momose, M. Inouye, Y. Maeda, H. Matsuzawa and T. Ohta. 1990. Enhancement of the thermostability of subtilisin E by introduction of a disulfide bond engineered on the basis of structural comparison with a thermophilic serine protease. J. Biolog.

Chem. 265(12):6874-6878.

Takami, H. and K. Horikoshi. 2000. Analysis of the genome of an alkaliphilic *Bacillus* strain from an industrial point of view. Extremophiles. 4:99-108.

Takami, H., K. Nakasone, N. Ogasawara, C. Hirama, Y. Nakamura, N. Masui, F. fuji, Y. Takaki, A. Inoue and K. Horikoshi. 1999. Sequencing of three lambda clones from the genome of alkaliphilic *Bacillus* sp. Strain C-125. Extremophiles 3: 29- 34.

Takami, H., S. Nakamura, R. Aono and K. Horikoshi. 1992. Degradation of human hair by a thermostable alkaline protease from alkaliphilic *Bacillus* sp. No. AH-101. Biosci. Biotechnol. Biochem. 56:1667-1669.

Takami, H., Y. Nogi and K. Horikoshi. 1999. Reidentification of the keratinase-producing facultatively alkaliphilic *Bacillus* sp. AH-101 as *Bacillus halodurans*. Extremophiles 3: 293- 296.

Tomarelli R. M., J. Charney and M. L. Harding. 1949. The use of azoalbumin as a substrate in the colorimetric determination of peptic and tryptic activity. J. Lab. Clin. Med. 34:428-433.

Williams, C. M., C. G. Lee, J. D. Garlich and J. C. H. Shih. 1990. Evaluation of a bacterial feather fermentation product, feather-lysate, as a feed protein. Poult. Sci. 70:85-94.

Williams, C. M., C. S. Richter, J. M. Mackenzie and J. C. H. Shih. 1990. Isolation, identification and characterication of a feather-degrading bacterium. Appl. Environ. Microbiol. 56(6): 1509-1515.