

Global Stability for the Leslie-Gower Predator-Prey System with Time-Delay and Holling's Type Functional Response

Han-Cheng Tsai and Chao-Pao Ho*

Abstract

In this paper, we concerned about the dynamical behavior of the Leslie-Gower predator-prey system with time delay and different functional response $p(x)$. At first, we discuss the global stability for the Leslie-Gower predator-prey system with time delay and $p(x) = cx$. Secondly, with $p(x) = \frac{cx}{1+x}$. Thirdly, with $p(x) = \frac{cx^2}{1+x^2}$. Finally, we illustrate our results by some examples.

1 Introduction

Predator-prey models have been studied for a long time. One of the most important problems in a predator-prey system is the global stability of the unique positive equilibrium point. The global stability analysis for predator-prey system without time delay has been done by many researchers. Most of them use the following methods to prove global stability of a predator-prey system without delay. The first method is to construct a Lyapunov function [3,4,5]. The second method is to employ the Dulac Criterion to eliminate the existence of periodic orbits and then use the Poincaré-Bendixson Theorem to analyze the global stability of the unique positive equilibrium [3,4,5,6]. The third method is the limit cycle stability analysis [3,6,7,8]. The fourth method is the comparison method [3,7,8].

But more realistic models should include some of the past states of the population system; that is, a real system should be modeled with time delays. In [9,10,11], authors were to analyze the global stability of the system with time delay by constructing a Lyapunov functional.

In this paper, we were concerned about the Leslie-Gower predator-prey system. For this system without delay as in [12], authors to analyze the global stability by constructing a Lyapunov

*Department of Mathematics, Tunghai University, Taichung 407, TAIWAN

function. And in [14], authors discussed the global stability of this system with a single delay by constructing a Lyapunov functional. Now, we are to establish global stability of the Leslie-Gower predator-prey system with a single delay with the different functional response of the predator $p(x)$, by constructing a Lyapunov functional. In section 2, we analyze the global stability of the Leslie-Gower predator-prey system with a single delay with $p(x) = cx$ in the Holling-type I, $p(x) = \frac{cx}{1+x}$ in the Holling-type II, and $p(x) = \frac{cx^2}{1+x^2}$ in the Holling-type III by constructing Lyapunov functionals. In section 3, we illustrate our results by some examples.

2 The Model with Time Delay

Consider the Leslie-Gower predator-prey system with time delay τ modeled by

$$\begin{aligned}\dot{x}(t) &= x(t) \left\{ r \left[1 - \frac{x(t-\tau)}{K} \right] \right\} - p(x(t))y(t) \\ \dot{y}(t) &= y(t) \left[\delta - \beta \frac{y(t)}{x(t)} \right]\end{aligned}\tag{2.1}$$

with the initial conditions

$$\begin{aligned}x(0) &= \phi(0) \geq 0, \quad \theta \in [-\tau, \infty), \quad \phi \in C^1([-\tau, \infty), \mathbb{R}) \\ x(0) &> 0, \quad y(0) > 0\end{aligned}\tag{2.2}$$

where δ, β, r , and τ are positive constants, K is defined as the prey environmental carrying capacity, x and y denote the densities of prey and predator population, respectively. Because all we want to discuss is biological population, we only consider the first quadrant in the $x - y$ plane. The following assumption is consistent with the system (2.1).

(A) $p \in C^1([0, \infty), [0, \infty))$; $p(0) = 0$ and $p'(x) \geq 0$ for all $x > 0$.

The functional response of the predator, $p(x)$, has been discussed in the literature. Now, we just concern about the following $p(x)$ of this paper, $p(x) = cx$ in the Holling-type I model, $p(x) = \frac{cx}{1+x}$ in the Holling-type II model, and $p(x) = \frac{cx^2}{1+x^2}$ in the Holling-type III model; where c is encounter rate.

Lemma 2.1 *Every solution of the system (2.1) with the initial conditions (2.2) exists in the interval $[0, \infty)$ and remains positive for all $t \geq 0$.*

Proof: It is true because

$$\begin{aligned}x(t) &= x(0) \exp \left\{ \int_0^t \left[r - \frac{rx(s-\tau)}{K} - \frac{p(x(s))}{x(s)} y(s) \right] ds \right\} \\y(t) &= y(0) \exp \left\{ \int_0^t \left[\delta - \beta \frac{y(s)}{x(s)} \right] ds \right\}\end{aligned}$$

and $x(0), y(0) > 0$.

Lemma 2.2 Let $(x(t), y(t))$ denote the solution of (2.1) with the initial conditions (2.2), then

$$0 < x(t) \leq M, 0 < y(t) \leq L \quad (2.3)$$

eventually for all large t , where

$$M = Ke^{r\tau} \quad (2.4)$$

$$L = \frac{\delta M}{\beta} \quad (2.5)$$

Proof: Now, we want to show that there exists a $T > 0$ such that $x(t) \leq M$ for $t > T$. By Lemma 2.1, we know that solutions of the system (2.1) are positive, and hence, by assumption (A), and (2.1)

$$\begin{aligned}\dot{x}(t) &= x(t) \left\{ r \left[1 - \frac{x(t-\tau)}{K} \right] \right\} - p(x(t))y(t) \\&\leq rx(t) \left[1 - \frac{x(t-\tau)}{K} \right]\end{aligned} \quad (2.6)$$

Taking $M^* = K(1 + K_1)$, $0 < K_1 < e^{r\tau} - 1$. Suppose $x(t)$ is not oscillatory about M^* . That is, there exists a $T > 0$ such that either

$$x(t) > M^* \quad \text{for } t > T_0 \quad (2.7)$$

or

$$x(t) \leq M^* \quad \text{for } t > T_0 \quad (2.8)$$

If (2.8) holds, then for $t > T_0$

$$x(t) \leq M^* = K(1 + K_1) < Ke^{r\tau} = M$$

That is, (2.3) holds. Suppose (2.7) holds. Equation (2.6) implies that for $t > T_0 + \tau$

$$\begin{aligned}\dot{x}(t) &\leq rx(t) \left[1 - \frac{x(t-\tau)}{K} \right] \\&< -K_1 rx(t)\end{aligned}$$

It follows that

$$\begin{aligned} \int_{T_0+\tau}^t \frac{\dot{x}(s)}{x(s)} ds &< \int_{T_0+\tau}^t -K_1 r ds \\ &= -K_1 r (t - T_0 - \tau) \end{aligned}$$

Then $0 < x(t) < x(T_0 + \tau)e^{-K_1 r(t - T_0 - \tau)} \rightarrow 0$ as $t \rightarrow \infty$. That is, $\lim_{t \rightarrow \infty} x(t) = 0$ by the Squeeze Theorem. It contradicts to (2.7). Therefore, there must exist a $T_1 > T_0$ such that $x(T_1) \leq M^*$. If $x(T_1) \leq M^*$ for all $t \geq T_1$, then (2.3) follows. If not, then there must exist a $T_2 > T_1$ such that T_2 be the first time which $x(T_2) > M^*$. Therefore, there exists a $T_3 > T_2$ such that T_3 be the first time which $x(T_3) < M^*$ by above discussion. By above, we know that $x(T_1) \leq M^*$, $x(T_2) > M^*$, and $x(T_3) \leq M^*$ where $T_1 < T_2 < T_3$. Then, by the Intermediate Value Theorem, there exists T_4 and T_5 such that

$$\begin{aligned} x(T_4) &= M^* , \quad T_1 \leq T_4 < T_2 \\ x(T_5) &= M^* , \quad T_2 \leq T_5 < T_3 \end{aligned}$$

and $x(t) > M^*$ for $T_4 < t < T_5$. Hence there is a $T_6 \in (T_4, T_5)$ such that $x(T_6)$ is an arbitrary local maximum, and hence it follows from (2.6) that

$$0 = \dot{x}(T_6) \leq rx(T_6) \left[1 - \frac{x(T_6 - \tau)}{K} \right]$$

and this implies

$$x(T_6 - \tau) \leq K$$

Integrating both sides of (2.6) on the interval $[T_6 - \tau, T_6]$, we have

$$\ln \left[\frac{x(T_6)}{x(T_6 - \tau)} \right] = \int_{T_6 - \tau}^{T_6} \frac{\dot{x}(s)}{x(s)} ds \leq \int_{T_6 - \tau}^{T_6} r \left[1 - \frac{x(s - \tau)}{K} \right] ds \leq r\tau$$

It follows that

$$x(T_6) \leq x(T_6 - \tau)e^{r\tau} \leq Ke^{r\tau} = M$$

Since $x(T_6)$ is local maximum of $x(t)$ and $x(T_6) \leq M$, $x(t) \leq M$ where t near T_6 . Since $x(T_6)$ is an arbitrary local maximum of $x(t)$, we can conclude that there exists a $T > 0$ such that

$$x(t) \leq M \quad \text{for } t \geq T \tag{2.9}$$

Suppose $x(t)$ is oscillatory about M^* , for this case, the proof is similarly to above one. Now, we want to show that $y(t)$ is bounded above by L eventually for all large t . By (2.9), it follows that for

$t > T$

$$\begin{aligned}\dot{y}(t) &= y(t) \left[\delta - \beta \frac{y(t)}{x(t)} \right] \\ &\leq y(t) \left[\delta - \frac{\beta}{M} y(t) \right] \\ &= \delta y(t) \left[1 - \frac{\beta}{\delta M} y(t) \right] \\ &= \delta y(t) \left[1 - \frac{y(t)}{\frac{\delta M}{\beta}} \right]\end{aligned}$$

Therefore, $y(t) \leq \frac{\delta M}{\beta} = L$ for $t > T$. This completes the proof.

Lemma 2.3 Suppose that the system (2.1) satisfies

$$r - cL > 0 \quad (2.10)$$

where L defined by (2.5), and c is defined in assumption (A). Then the system (2.1) is uniformly persistent. That is, there exists m, l , and $T^* > 0$ such that $m \leq x \leq M$ and $l \leq y \leq L$ for $t \geq T^*$.

Proof: By Lemma 2.2, and assumption (A), equation (2.1) follows that for $t \geq T + \tau$

$$\dot{x}(t) \geq x(t) \left[r \left(1 - \frac{M}{K} \right) - cL \right] \quad (2.11)$$

Integrating both sides of (2.11) on $[t - \tau, t]$, where $t \geq T + \tau$, then we have

$$x(t) \geq x(t - \tau) e^{[r(1 - \frac{M}{K}) - cL]\tau}$$

That is

$$x(t - \tau) \leq x(t) e^{-[r(1 - \frac{M}{K}) - cL]\tau} \quad (2.12)$$

It follows from (2.1) that for $t \geq T + \tau$

$$\begin{aligned}\dot{x}(t) &= x(t) r \left[1 - \frac{x(t - \tau)}{K} \right] - p(x(t)) y(t) \\ &\geq x(t) \left\{ r - cL - \frac{r}{K} e^{-[r(1 - \frac{M}{K}) - cL]\tau} x(t) \right\} \\ &= (r - cL) x(t) \left\{ 1 - \frac{x(t)}{\frac{K(r - cL)}{r} e^{[r(1 - \frac{M}{K}) - cL]\tau}} \right\}\end{aligned}$$

It follows that

$$\liminf_{t \rightarrow \infty} x(t) \geq \frac{K(r - cL)}{r} e^{[r(1 - \frac{M}{K}) - cL]\tau} \equiv \bar{m}$$

and $\bar{m} > 0$ by (2.10). So, for large t , $x(t) > \frac{\bar{m}}{2} \equiv m > 0$. It follows that

$$\begin{aligned}\dot{y}(t) &\geq y(t) \left[\delta - \beta \frac{y(t)}{m} \right] \\ &= \delta y(t) \left[1 - \frac{\beta}{\delta m} y(t) \right] \\ &= \delta y(t) \left[1 - \frac{y(t)}{\frac{\delta m}{\beta}} \right]\end{aligned}$$

Then

$$\liminf_{t \rightarrow \infty} y(t) \geq \frac{\delta m}{\beta} \equiv \bar{l}$$

So, for large t , $y(t) > \frac{\bar{l}}{2} \equiv l > 0$. Let

$$D = \{(x, y) | m \leq x \leq M, l \leq y \leq L\}$$

Then D is bounded compact region in R_+^2 that has positive distance from coordinate hyperplanes. Hence we obtain that there exists a $T^* > 0$ such that if $t \geq T^*$, then every positive solution of system (2.1) with the initial conditions (2.2) eventually enters and remains in the region D , that is, system (2.1) is uniformly persistent.

Theorem 2.1 If $p(x) = cx$ in the Holling-type I model, and the delay τ satisfy

$$r - cL > 0 \quad (2.13)$$

$$\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mr^2x^*\tau}{K^2} - \frac{Mrcy^*\tau}{2K} > 0 \quad (2.14)$$

$$\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mrcy^*\tau}{2K} > 0 \quad (2.15)$$

where m , M , and L defined in Lemmas 2.2 and 2.3, then the unique positive equilibrium E^* of the system (2.1) is globally asymptotically stable.

Proof: Define $z(t) = (z_1(t), z_2(t))$ by

$$z_1(t) = \frac{x(t) - x^*}{x^*}, \quad z_2(t) = \frac{y(t) - y^*}{y^*}$$

From (2.1),

$$\dot{z}_1(t) = [1 + z_1(t)] \left[-\frac{rx^*}{K} z_1(t - \tau) - cy^* z_2(t) \right] \quad (2.16)$$

$$\dot{z}_2(t) = [1 + z_2(t)] \left\{ \frac{\delta x^* z_1(t) - \beta y^* z_2(t)}{x^* [1 + z_1(t)]} \right\} \quad (2.17)$$

Let

$$V_1(z(t)) = \{z_1(t) - \ln[1 + z_1(t)]\} + \{z_2(t) - \ln[1 + z_2(t)]\} \quad (2.18)$$

then we have from (2.16) and (2.17) that

$$\begin{aligned} \dot{V}_1(z(t)) &= \frac{z_1(t)\dot{z}_1(t)}{1+z_1(t)} + \frac{z_2(t)\dot{z}_2(t)}{1+z_2(t)} \\ &= -\frac{rx^*}{K}z_1(t)z_1(t-\tau) - cy^*z_1(t)z_2(t) + \frac{\delta}{1+z_1(t)}z_1(t)z_2(t) \\ &\quad - \frac{\beta y^* z_2^2(t)}{x^*[1+z_1(t)]} \\ &\leq -\frac{rx^*}{K}z_1(t)z_1(t-\tau) + \frac{\delta - cy^*[1+z_1(t)]}{1+z_1(t)}z_1(t)z_2(t) \\ &\quad - \frac{\beta y^* z_2^2(t)}{x^*[1+z_1(t)]} \end{aligned} \quad (2.19)$$

If $\delta x^* - cy^*M > 0$, and by Lemma 2.3, there exists a $T^* > 0$ such that $m \leq x^*[1 + z_1(t)] \leq M$ and $l \leq y^*[1 + z_2(t)] \leq L$ for $t > T^*$. Then (2.19) implies that

$$\begin{aligned} \dot{V}_1(z(t)) &\leq -\frac{rx^*}{K}z_1(t)z_1(t-\tau) \\ &\quad + \frac{\delta x^* - cy^*m}{2m}[z_1^2(t) + z_2^2(t)] - \frac{\beta y^* z_2^2(t)}{M} \\ &= -\frac{rx^*}{K}z_1(t) \left[z_1(t) - \int_{t-\tau}^t \dot{z}_1(s)ds \right] + \left(\frac{\delta x^*}{2m} - \frac{cy^*}{2} \right) z_1^2(t) \\ &\quad + \left(\frac{\delta x^*}{2m} - \frac{cy^*}{2} - \frac{\beta y^*}{M} \right) z_2^2(t) \\ &= -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} \right) z_1^2(t) - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} \right) z_2^2(t) \\ &\quad + \frac{rx^*}{K}z_1(t) \int_{t-\tau}^t [1 + z_1(s)] \left[-\frac{rx^*}{K}z_1(s-\tau) - cy^*z_2(s) \right] ds \\ &= -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} \right) z_1^2(t) - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} \right) z_2^2(t) \\ &\quad + \frac{rx^*}{K} \int_{t-\tau}^t [1 + z_1(s)] \left[-\frac{rx^*}{K}z_1(t)z_1(s-\tau) - cy^*z_1(t)z_2(s) \right] ds \\ &\leq -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} \right) z_1^2(t) - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} \right) z_2^2(t) \\ &\quad + \frac{rx^*}{K} \int_{t-\tau}^t [1 + z_1(s)] \left[\frac{rx^*}{K}|z_1(t)||z_1(s-\tau)| + cy^*|z_1(t)||z_2(s)| \right] ds \end{aligned} \quad (2.20)$$

Then for $t \geq T^* + \tau \equiv \hat{T}$, we have from (2.20) that

$$\begin{aligned}
\dot{V}_1(z(t)) &\leq -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m}\right)z_1^2(t) - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m}\right)z_2^2(t) \\
&\quad + \frac{Mr}{K} \int_{t-\tau}^t \left[\frac{rx^*}{K} |z_1(t)| |z_1(s-\tau)| + cy^* |z_1(t)| |z_2(s)| \right] ds \\
&\leq -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m}\right)z_1^2(t) - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m}\right)z_2^2(t) \\
&\quad + \frac{Mr}{K} \left[\frac{rx^*\tau}{2K} z_1^2(t) + \frac{rx^*}{2K} \int_{t-\tau}^t z_1^2(s-\tau) ds + \frac{cy^*\tau}{2} z_1^2(t) \right. \\
&\quad \left. + \frac{cy^*}{2} \int_{t-\tau}^t z_2^2(s) ds \right] \\
&= -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mr^2x^*\tau}{2K^2} - \frac{Mrcy^*\tau}{2K}\right)z_1^2(t) - \left(\frac{\beta y^*}{M} \right. \\
&\quad \left. + \frac{cy^*}{2} - \frac{\delta x^*}{2m}\right)z_2^2(t) + \frac{Mr^2x^*}{2K^2} \int_{t-\tau}^t z_1^2(s-\tau) ds + \frac{Mrcy^*}{2K} \int_{t-\tau}^t z_2^2(s) ds
\end{aligned} \tag{2.21}$$

Let

$$\begin{aligned}
V_2(z(t)) &= \frac{Mr^2x^*}{2K^2} \int_{t-\tau}^t \int_s^t z_1^2(\gamma-\tau) d\gamma ds \\
&\quad + \frac{Mrcy^*}{2K} \int_{t-\tau}^t \int_s^t z_2^2(\gamma) d\gamma ds
\end{aligned} \tag{2.22}$$

then

$$\begin{aligned}
\dot{V}_2(z(t)) &= \frac{Mr^2x^*\tau}{2K^2} z_1^2(t-\tau) - \frac{Mr^2x^*}{2K^2} \int_{t-\tau}^t z_1^2(s-\tau) ds \\
&\quad + \frac{Mrcy^*\tau}{2K} z_2^2(t) - \frac{Mrcy^*}{2K} \int_{t-\tau}^t z_2^2(s) ds
\end{aligned} \tag{2.23}$$

and then we have from (2.21) and (2.23) that for $t \geq \hat{T}$

$$\begin{aligned}
\dot{V}_1(z(t)) + \dot{V}_2(z(t)) &\leq -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mr^2x^*\tau}{2K^2} - \frac{Mrcy^*\tau}{2K}\right)z_1^2(t) \\
&\quad - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mrcy^*\tau}{2K}\right)z_2^2(t) \\
&\quad + \frac{Mr^2x^*\tau}{2K^2} z_1^2(t-\tau)
\end{aligned} \tag{2.24}$$

Let

$$V_3(z(t)) = \frac{Mr^2x^*\tau}{2K^2} \int_{t-\tau}^t z_1^2(s) ds \tag{2.25}$$

then

$$\dot{V}_3(z(t)) = \frac{Mr^2x^*\tau}{2K^2}z_1^2(t) - \frac{Mr^2x^*\tau}{2K^2}z_1^2(t-\tau) \quad (2.26)$$

Now define a Lyapunov functional $V(z(t))$ as

$$V(z(t)) = V_1(z(t)) + V_2(z(t)) + V_3(z(t)) \quad (2.27)$$

then we have from (2.24) and (2.26) that for $t \geq \hat{T}$

$$\begin{aligned} \dot{V}(z(t)) &\leq -\left(\frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mr^2x^*\tau}{K^2} - \frac{Mrcy^*\tau}{2K}\right)z_1^2(t) \\ &\quad - \left(\frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mrcy^*\tau}{2K}\right)z_2^2(t) \\ &\equiv -\zeta z_1^2(t) - \eta z_2^2(t) \end{aligned} \quad (2.28)$$

Then it follows from (2.14) and (2.15) that $\zeta > 0$ and $\eta > 0$. Let $w(s) = \hat{N}s^2$ where $\hat{N} = \min\{\zeta, \eta\}$, then w is nonnegative continuous on $[0, \infty)$, $w(0) = 0$, and $w(s) > 0$ for $s > 0$. It follows from (2.28) that for $t \geq \hat{T}$

$$\dot{V}(z(t)) \leq -\hat{N}[z_1^2(t) + z_2^2(t)] = -\hat{N}\|z(t)\|^2 = -w(\|z(t)\|) \quad (2.29)$$

Now, we want to find a function u such that $V(z(t)) \geq u(\|z(t)\|)$. It follows from (2.18), (2.22), and (2.25) that

$$V(z(t)) \geq \{z_1(t) - \ln[1 + z_1(t)]\} + \{z_2(t) - \ln[1 + z_2(t)]\} \quad (2.30)$$

By the Taylor Theorem, we have that

$$z_i(t) - \ln[1 + z_i(t)] = \frac{z_i^2(t)}{2[1 + \theta_i(t)]^2} \quad (2.31)$$

where $\theta_i(t) \in (0, z_i(t))$ or $(z_i(t), 0)$ for $i = 1, 2$.

Case 1 : If $0 < \theta_i(t) < z_i(t)$ for $i = 1, 2$, then

$$\frac{z_i^2(t)}{[1 + z_i(t)]^2} < \frac{z_i^2(t)}{[1 + \theta_i(t)]^2} < z_i^2(t) \quad (2.32)$$

By Lemma 2.3, it follows that for $t \geq T^*$

$$\begin{aligned} m \leq x^*[1 + z_1(t)] &= x(t) \leq M \\ l \leq y^*[1 + z_2(t)] &= y(t) \leq L \end{aligned} \quad (2.33)$$

Then (2.32) implies that

$$\begin{aligned} \left(\frac{x^*}{M}\right)^2 z_1^2(t) &\leq \frac{z_1^2(t)}{[1+\theta_1(t)]^2} < z_1^2(t) \\ \left(\frac{y^*}{L}\right)^2 z_2^2(t) &\leq \frac{z_2^2(t)}{[1+\theta_2(t)]^2} < z_2^2(t) \end{aligned} \quad (2.34)$$

It follows that (2.30), (2.31), and (2.34) that for $t \geq T^*$

$$\begin{aligned} V(z(t)) &\geq \frac{1}{2} \frac{z_1^2(t)}{[1+\theta_1(t)]^2} + \frac{1}{2} \frac{z_2^2(t)}{[1+\theta_2(t)]^2} \\ &\geq \frac{1}{2} \left(\frac{x^*}{M}\right)^2 z_1^2(t) + \frac{1}{2} \left(\frac{y^*}{L}\right)^2 z_2^2(t) \\ &\geq \min \left\{ \frac{1}{2} \left(\frac{x^*}{M}\right)^2, \frac{1}{2} \left(\frac{y^*}{L}\right)^2 \right\} \cdot [z_1^2(t) + z_2^2(t)] \\ &\equiv \tilde{N} \|z(t)\|^2 \end{aligned}$$

Case2 : If $-1 < z_i(t) < \theta_i(t) < 0$ for $i = 1, 2$, then

$$z_i^2(t) < \frac{z_i^2(t)}{[1+\theta_i(t)]^2} < \frac{z_i^2(t)}{[1+z_i(t)]^2} \quad (2.35)$$

By (2.33), (2.35) implies that

$$\begin{aligned} z_1^2(t) &< \frac{z_1^2(t)}{[1+\theta_1(t)]^2} \leq \left(\frac{x^*}{m}\right)^2 z_1^2(t) \\ z_2^2(t) &< \frac{z_2^2(t)}{[1+\theta_2(t)]^2} \leq \left(\frac{y^*}{l}\right)^2 z_2^2(t) \end{aligned} \quad (2.36)$$

It follows that (2.30), (2.31), and (2.36) that for $t \geq T^*$

$$\begin{aligned} V(z(t)) &\geq \frac{1}{2} \frac{z_1^2(t)}{[1+\theta_1(t)]^2} + \frac{1}{2} \frac{z_2^2(t)}{[1+\theta_2(t)]^2} \\ &> \frac{1}{2} z_1^2(t) + \frac{1}{2} z_2^2(t) \\ &\geq \frac{1}{2} \left(\frac{x^*}{M}\right)^2 z_1^2(t) + \frac{1}{2} \left(\frac{y^*}{L}\right)^2 z_2^2(t) \\ &\geq \tilde{N} [z_1^2(t) + z_2^2(t)] \\ &= \tilde{N} \|z(t)\|^2 \end{aligned}$$

Case3 : If $0 < \theta_1(t) < z_1(t)$ and $-1 < z_2(t) < \theta_2(t) < 0$, then it follows that (2.30), (2.31), (2.34), and (2.36) that for $t \geq T^*$

$$\begin{aligned} V(z(t)) &\geq \frac{1}{2} \frac{z_1^2(t)}{[1 + \theta_1(t)]^2} + \frac{1}{2} \frac{z_2^2(t)}{[1 + \theta_2(t)]^2} \\ &> \frac{1}{2} \left(\frac{x^*}{M} \right)^2 z_1^2(t) + \frac{1}{2} z_2^2(t) \\ &\geq \frac{1}{2} \left(\frac{x^*}{M} \right)^2 z_1^2(t) + \frac{1}{2} \left(\frac{y^*}{L} \right)^2 z_2^2(t) \\ &\geq \tilde{N} [z_1^2(t) + z_2^2(t)] \\ &= \tilde{N} \|z(t)\|^2 \end{aligned}$$

Case4 : If $-1 < z_1(t) < \theta_1(t) < 0$ and $0 < \theta_2(t) < z_2(t)$, then it follows that (2.30), (2.31), (2.34), and (2.36) that for $t \geq T^*$

$$\begin{aligned} V(y(t)) &\geq \frac{1}{2} \frac{z_1^2(t)}{[1 + \theta_1(t)]^2} + \frac{1}{2} \frac{z_2^2(t)}{[1 + \theta_2(t)]^2} \\ &> \frac{1}{2} z_1^2(t) + \frac{1}{2} \left(\frac{y^*}{L} \right)^2 z_2^2(t) \\ &\geq \frac{1}{2} \left(\frac{x^*}{M} \right)^2 z_1^2(t) + \frac{1}{2} \left(\frac{y^*}{L} \right)^2 z_2^2(t) \\ &\geq \tilde{N} [z_1^2(t) + z_2^2(t)] \\ &= \tilde{N} \|z(t)\|^2 \end{aligned}$$

Let $u(s) = \tilde{N}s^2$, then u is nonnegative continuous on $[0, \infty)$, $u(0) = 0$, $u(s) > 0$ for $s > 0$, and $\lim_{s \rightarrow \infty} u(s) = +\infty$. So, by case1 ~ case4, we have

$$V(z(t)) \geq u(\|z(t)\|) \quad \text{for } t \geq T^* \quad (2.37)$$

So the equilibrium point E^* of the system (2.1) is globally asymptotically stable with $p(x) = cx$.

Theorem 2.2 If $p(x) = \frac{cx}{1+x}$ in the Holling-type II model, and the delay τ satisfy

$$r - cL > 0 \quad (2.38)$$

$$\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} - \frac{rx^*(x^*+M)}{K(1+m)}$$

$$-\frac{Mr^2x^*\tau(1+x^*)(K+3x^*+1+M)}{K^2(1+m)^2} - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} > 0 \quad (2.39)$$

$$\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} > 0 \quad (2.40)$$

where m , M , and L defined in Lemmas 2.2 and 2.3, then the unique positive equilibrium E^* of the system (2.1) is globally asymptotically stable.

Proof: Define $z(t) = (z_1(t), z_2(t))$ by

$$z_1(t) = \frac{x(t) - x^*}{x^*}, \quad z_2(t) = \frac{y(t) - y^*}{y^*}$$

From (2.1),

$$\begin{aligned} \dot{z}_1(t) &= [1+z_1(t)] \left\{ \frac{rx^*z_1(t)}{1+x^*[1+z_1(t)]} - \frac{rx^{*2}z_1(t)}{K\{1+x^*[1+z_1(t)]\}} \right. \\ &\quad - \frac{rx^*z_1(t-\tau)}{K\{1+x^*[1+z_1(t)]\}} - \frac{rx^{*2}z_1(t-\tau)}{K\{1+x^*[1+z_1(t)]\}} \\ &\quad \left. - \frac{rx^{*2}z_1(t-\tau)z_1(t)}{K\{1+x^*[1+z_1(t)]\}} - \frac{cy^*z_2(t)}{1+x^*[1+z_1(t)]} \right\} \quad (2.41) \\ \dot{z}_2(t) &= [1+z_2(t)] \left\{ \frac{\delta x^*z_1(t) - \beta y^*z_2(t)}{x^*[1+z_1(t)]} \right\} \quad (2.42) \end{aligned}$$

Let

$$V_1(z(t)) = \{z_1(t) - \ln[1+z_1(t)]\} + \{z_2(t) - \ln[1+z_2(t)]\} \quad (2.43)$$

then we have from (2.41) and (2.42) that

$$\begin{aligned} \dot{V}_1(z(t)) &= \frac{\dot{z}_1(t)z_1(t)}{1+z_1(t)} + \frac{\dot{z}_2(t)z_2(t)}{1+z_2(t)} \\ &= \frac{rx^*z_1^2(t)}{1+x^*[1+z_1(t)]} - \frac{rx^{*2}z_1^2(t)}{K\{1+x^*[1+z_1(t)]\}} - \frac{rx^*z_1(t)z_1(t-\tau)}{K\{1+x^*[1+z_1(t)]\}} \\ &\quad - \frac{rx^{*2}z_1(t)z_1(t-\tau)}{K\{1+x^*[1+z_1(t)]\}} - \frac{rx^{*2}z_1^2(t)z_1(t-\tau)}{K\{1+x^*[1+z_1(t)]\}} - \frac{\beta y^*z_2^2(t)}{x^*[1+z_1(t)]} \\ &\quad + \frac{\delta\{1+x^*[1+z_1(t)]\} - cy^*[1+z_1(t)]}{\{1+x^*[1+z_1(t)]\}[1+z_1(t)]} z_1(t)z_2(t) \\ &\leq \frac{rx^*z_1^2(t)}{1+x^*[1+z_1(t)]} - \frac{rx^{*2}z_1^2(t)}{K\{1+x^*[1+z_1(t)]\}} + \frac{rx^{*2}z_1^2(t)|z_1(t-\tau)|}{K\{1+x^*[1+z_1(t)]\}} \\ &\quad - \frac{\beta y^*z_2^2(t)}{x^*[1+z_1(t)]} + \frac{\delta\{1+x^*[1+z_1(t)]\} - cy^*[1+z_1(t)]}{\{1+x^*[1+z_1(t)]\}[1+z_1(t)]} z_1(t)z_2(t) \\ &\quad - \left\{ \frac{rx^*}{K\{1+x^*[1+z_1(t)]\}} + \frac{rx^{*2}}{K\{1+x^*[1+z_1(t)]\}} \right\} z_1(t)z_1(t-\tau) \quad (2.44) \end{aligned}$$

If $\delta x^*(1+m) - cy^*M > 0$, and by Lemma 2.3, there exists a $T^* > 0$ such that $m \leq x^*[1+z_1(t)] \leq M$ and $l \leq y^*[1+z_2(t)] \leq L$ for $t > T^*$. Then (2.44) implies that

$$\begin{aligned}
\dot{V}_1(z(t)) &\leq \frac{rx^*z_1^2(t)}{1+m} - \frac{rx^{*2}z_1^2(t)}{K(1+M)} + \frac{rx^{*2}z_1^2(t)|z_1(t-\tau)|}{K(1+m)} \\
&\quad - \frac{\beta y^*z_2^2(t)}{M} - \frac{cy^*z_1^2(t)}{2(1+m)} - \frac{cy^*z_2^2(t)}{2(1+m)} + \frac{\delta x^*(1+M)z_1^2(t)}{2m(1+m)} \\
&\quad + \frac{\delta x^*(1+M)z_2^2(t)}{2m(1+m)} - \frac{rx^*(1+x^*)}{K\{1+x^*[1+z_1(t)]\}} z_1(t) \left[z_1(t) - \int_{t-\tau}^t \dot{z}_1(s) ds \right] \\
&= - \left[\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} \right] z_1^2(t) \\
&\quad - \left[\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} \right] z_2^2(t) + \frac{rx^{*2}}{K(1+m)} z_1^2(t) |z_1(t-\tau)| \\
&\quad + \frac{rx^*(1+x^*)}{K\{1+x^*[1+z_1(t)]\}} \int_{t-\tau}^t [1+z_1(s)] \left\{ \frac{rx^*z_1(t)z_1(s)}{1+x^*[1+z_1(s)]} \right. \\
&\quad \left. - \frac{rx^{*2}z_1(t)z_1(s)}{K\{1+x^*[1+z_1(s)]\}} - \frac{rx^*z_1(t)z_1(s-\tau)}{K\{1+x^*[1+z_1(s)]\}} - \frac{rx^{*2}z_1(t)z_1(s-\tau)}{K\{1+x^*[1+z_1(s)]\}} \right. \\
&\quad \left. - \frac{rx^{*2}z_1(t)z_1(s)z_1(s-\tau)}{K\{1+x^*[1+z_1(s)]\}} - \frac{cy^*z_1(t)z_2(s)}{1+x^*[1+z_1(s)]} \right\} ds \\
&\leq - \left[\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} \right] z_1^2(t) \\
&\quad - \left[\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} \right] z_2^2(t) + \frac{rx^{*2}}{K(1+m)} z_1^2(t) |z_1(t-\tau)| \\
&\quad + \frac{rx^*(1+x^*)}{K\{1+x^*[1+z_1(t)]\}} \int_{t-\tau}^t [1+z_1(s)] \left\{ \frac{rx^*|z_1(t)||z_1(s)|}{1+x^*[1+z_1(s)]} \right. \\
&\quad \left. + \frac{rx^{*2}|z_1(t)||z_1(s)|}{K\{1+x^*[1+z_1(s)]\}} + \frac{rx^*|z_1(t)||z_1(s-\tau)|}{K\{1+x^*[1+z_1(s)]\}} \right. \\
&\quad \left. + \frac{rx^{*2}|z_1(t)||z_1(s-\tau)|}{K\{1+x^*[1+z_1(s)]\}} + \frac{cy^*|z_1(t)||z_2(s)|}{1+x^*[1+z_1(s)]} \right\} ds \\
&\quad - \frac{rx^*(1+x^*)}{K\{1+x^*[1+z_1(t)]\}} \int_{t-\tau}^t \frac{rx^{*2}z_1(t)z_1(s-\tau)}{K\{1+x^*[1+z_1(s)]\}} [1+z_1(s)]^2 ds \\
&\quad + \frac{rx^*(1+x^*)}{K\{1+x^*[1+z_1(t)]\}} \int_{t-\tau}^t \frac{rx^{*2}|z_1(t)||z_1(s-\tau)|}{K\{1+x^*[1+z_1(s)]\}} [1+z_1(s)] ds \tag{2.45}
\end{aligned}$$

Then for $t \geq T^* + \tau \equiv \hat{T}$, we have from (2.45) that

$$\begin{aligned}
\dot{V}_1(z(t)) &\leq - \left[\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} \right] z_1^2(t) \\
&\quad - \left[\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} \right] z_2^2(t) + \frac{rx^{*2}}{K(1+m)} \left(1 + \frac{M}{x^*} \right) z_1^2(t) \\
&\quad + \frac{Mr(1+x^*)}{K(1+m)} \int_{t-\tau}^t \left[\frac{rx^*|z_1(t)||z_1(s)|}{1+m} + \frac{rx^{*2}|z_1(t)||z_1(s)|}{K(1+m)} \right. \\
&\quad \left. + \frac{rx^*|z_1(t)||z_1(s-\tau)|}{K(1+m)} + \frac{rx^{*2}|z_1(t)||z_1(s-\tau)|}{K(1+m)} + \frac{cy^*|z_1(t)||z_2(s)|}{1+m} \right] ds \\
&\quad + \frac{M^2r(1+x^*)}{Kx^*(1+m)} \int_{t-\tau}^t \frac{rx^{*2}|z_1(t)||z_1(s-\tau)|}{K(1+m)} ds \\
&\quad + \frac{Mr(1+x^*)}{K(1+m)} \int_{t-\tau}^t \frac{rx^{*2}|z_1(t)||z_1(s-\tau)|}{K(1+m)} ds \\
&\leq - \left[\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} \right. \\
&\quad \left. - \frac{rx^*(x^*+M)}{K(1+m)} - \frac{Mr(1+x^*)\tau}{K(1+m)^2} \left(\frac{rx^*}{2} + \frac{3rx^{*2}}{2K} + \frac{rx^*}{2K} + \frac{cy^*}{2} \right) \right. \\
&\quad \left. - \frac{M^2r^2x^*(1+x^*)\tau}{2K^2(1+m)^2} \right] z_1^2(t) - \left[\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*}{2m} \right] z_2^2(t) \\
&\quad + \frac{Mr^2x^*(1+x^*)(K+x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t z_1^2(s) ds \\
&\quad + \frac{Mr^2x^*(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t z_1^2(s-\tau) ds \\
&\quad + \frac{Mrcy^*(1+x^*)}{2K(1+m)^2} \int_{t-\tau}^t z_2^2(s) ds
\end{aligned} \tag{2.46}$$

Let

$$\begin{aligned}
V_2(z(t)) &= \frac{Mr^2x^*(1+x^*)(K+x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t \int_s^t z_1^2(\gamma) d\gamma ds \\
&\quad + \frac{Mr^2x^*(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t \int_s^t z_1^2(\gamma-\tau) d\gamma ds \\
&\quad + \frac{Mrcy^*(1+x^*)}{2K(1+m)^2} \int_{t-\tau}^t \int_s^t z_2^2(\gamma) d\gamma ds
\end{aligned} \tag{2.47}$$

then

$$\begin{aligned}
\dot{V}_2(z(t)) &= \frac{Mr^2x^*\tau(1+x^*)(K+x^*)}{2K^2(1+m)^2}z_1^2(t) \\
&\quad - \frac{Mr^2x^*(1+x^*)(K+x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t z_1^2(s)ds \\
&\quad + \frac{Mr^2x^*\tau(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2}z_1^2(t-\tau) \\
&\quad - \frac{Mr^2x^*(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t z_1^2(s-\tau)ds \\
&\quad + \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2}z_2^2(t) - \frac{Mrcy^*(1+x^*)}{2K(1+m)^2} \int_{t-\tau}^t z_2^2(s)ds
\end{aligned} \tag{2.48}$$

and then we have from (2.46) and (2.48) that for $t \geq \hat{T}$

$$\begin{aligned}
\dot{V}_1(z(t)) + \dot{V}_2(z(t)) &\leq - \left[\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} \right. \\
&\quad - \frac{rx^*(x^*+M)}{K(1+m)} - \frac{Mr(1+x^*)\tau}{K(1+m)^2} \left(\frac{rx^*}{2} + \frac{3rx^{*2}}{2K} + \frac{rx^*}{2K} + \frac{cy^*}{2} \right) \\
&\quad \left. - \frac{M^2r^2x^*\tau(1+x^*)}{2K^2(1+m)^2} - \frac{Mr^2x^*\tau(1+x^*)(K+x^*)}{2K^2(1+m)^2} \right] z_1^2(t) \\
&\quad - \left[\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} \right] z_2^2(t) \\
&\quad + \frac{Mr^2x^*\tau(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} z_1^2(t-\tau)
\end{aligned} \tag{2.49}$$

Let

$$V_3(z(t)) = \frac{Mr^2x^*\tau(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} \int_{t-\tau}^t z_1^2(s)ds \tag{2.50}$$

then

$$\begin{aligned}
\dot{V}_3(z(t)) &= \frac{Mr^2x^*\tau(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} z_1^2(t) \\
&\quad - \frac{Mr^2x^*\tau(1+x^*)(M+1+2x^*)}{2K^2(1+m)^2} z_1^2(t-\tau)
\end{aligned} \tag{2.51}$$

Now define a Lyapunov functional $V(z(t))$ as

$$V(z(t)) = V_1(z(t)) + V_2(z(t)) + V_3(z(t)) \quad (2.52)$$

then we have from (2.49) and (2.51) that for $t \geq \hat{T}$

$$\begin{aligned} \dot{V}(z(t)) &\leq -\left[\frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} \right. \\ &\quad \left. - \frac{rx^*(x^*+M)}{K(1+m)} - \frac{Mr^2x^*\tau(1+x^*)(K+3x^*+1+M)}{K^2(1+m)^2} \right. \\ &\quad \left. - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} \right] z_1^2(t) \\ &\quad - \left[\frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} \right] z_2^2(t) \\ &\equiv -\zeta z_1^2(t) - \eta z_2^2(t) \end{aligned} \quad (2.53)$$

Then it follows from (2.39) and (2.40) that $\zeta > 0$ and $\eta > 0$. Let $w(s) = \hat{N}s^2$ where $\hat{N} = \min\{\zeta, \eta\}$, then w is nonnegative continuous on $[0, \infty)$, $w(0) = 0$, and $w(s) > 0$ for $s > 0$. It follows from (2.53) that for $t \geq \hat{T}$

$$\dot{V}(z(t)) \leq -\hat{N}[z_1^2(t) + z_2^2(t)] = -\hat{N}\|z(t)\|^2 = -w(\|z(t)\|) \quad (2.54)$$

Now, we want to find a function u such that $V(z(t)) \geq u(\|z(t)\|)$. It follows from (2.43), (2.47), and (2.50) that

$$V(z(t)) \geq \{z_1(t) - \ln[1+z_1(t)]\} + \{z_2(t) - \ln[1+z_2(t)]\} \quad (2.55)$$

then by Theorem 2.1, we have that $u(s) = \tilde{N}s^2$ and $V(z(t)) \geq u(\|z(t)\|)$. So the equilibrium point E^* of the system (2.1) is globally asymptotically stable with $p(x) = \frac{cx}{1+x}$.

Theorem 2.3 If $p(x) = \frac{cx^2}{1+x^2}$ in the Holling-type III model, and the delay τ satisfy

$$r - cL > 0 \quad (2.56)$$

$$\begin{aligned}
& \frac{rx^*(1+3x^{*2})}{K(1+M^2)} + \frac{cx^*y^*}{2(1+m^2)} + \frac{cx^*y^*}{1+M^2} - \frac{rx^*(M+3x^*)}{1+m^2} - \frac{3rx^{*2}(M+x^*)}{K(1+m^2)} \\
& - \frac{rx^*(M+x^*)^2}{K(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} - \frac{cx^*(L+y^*)}{1+m^2} \\
& - \frac{Mr^2x^*\tau(1+x^{*2})(3Kx^*+9x^{*2}+MK+5Mx^*+1+M^2)}{K^2(1+m^2)^2} \\
& - \frac{Mrcy^*\tau(1+x^{*2})(4x^*+M)}{2K(1+m^2)^2} > 0
\end{aligned} \tag{2.57}$$

$$\frac{\beta y^*}{M} + \frac{cx^*y^*}{2(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} - \frac{Mrcy^*\tau(1+x^{*2})(2x^*+M)}{2K(1+m^2)^2} > 0 \tag{2.58}$$

where m , M , and L defined in Lemmas 2.2 and 2.3, then the unique positive equilibrium E^* of the system (2.1) is globally asymptotically stable.

Proof: Define $z(t) = (z_1(t), z_2(t))$ by

$$z_1(t) = \frac{x(t) - x^*}{x^*}, \quad z_2(t) = \frac{y(t) - y^*}{y^*}$$

From (2.1),

$$\begin{aligned}
\dot{z}_1(t) = & [1+z_1(t)] \left\{ \frac{2rx^{*2}z_1(t)}{1+x^{*2}[1+z_1(t)]^2} + \frac{rx^{*2}z_1^2(t)}{1+x^{*2}[1+z_1(t)]^2} \right. \\
& - \frac{2rx^{*3}z_1(t)}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{rx^{*3}z_1^2(t)}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\
& - \frac{rx^*z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{rx^*z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\
& - \frac{2rx^{*3}z_1(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{rx^{*3}z_1^2(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\
& \left. - \frac{cx^*y^*z_1(t)}{1+x^{*2}[1+z_1(t)]^2} - \frac{cx^*y^*z_2(t)}{1+x^{*2}[1+z_1(t)]^2} - \frac{cx^*y^*z_1(t)z_2(t)}{1+x^{*2}[1+z_1(t)]^2} \right\}
\end{aligned} \tag{2.59}$$

$$\dot{z}_2(t) = [1+z_2(t)] \left\{ \frac{\delta x^*z_1(t) - \beta y^*z_2(t)}{x^*[1+z_1(t)]} \right\} \tag{2.60}$$

Let

$$V_1(z(t)) = \{z_1(t) - \ln[1 + z_1(t)]\} + \{z_2(t) - \ln[1 + z_2(t)]\} \quad (2.61)$$

then we have from (2.59) and (2.60) that

$$\begin{aligned} \dot{V}_1(z(t)) &= \frac{\dot{z}_1(t)z_1(t)}{1+z_1(t)} + \frac{\dot{z}_2(t)z_2(t)}{1+z_2(t)} \\ &= \frac{2rx^{*2}z_1^2(t)}{1+x^{*2}[1+z_1(t)]^2} + \frac{rx^{*2}z_1^3(t)}{1+x^{*2}[1+z_1(t)]^2} \\ &\quad - \frac{2rx^{*3}z_1^2(t)}{K\{1+x_1^{*2}[1+z_1(t)]^2\}} - \frac{rx^{*3}z_1^3(t)}{K\{1+x_1^{*2}[1+z_1(t)]^2\}} \\ &\quad - \frac{rx^{*}z_1(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{rx^{*3}z_1(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\ &\quad - \frac{2rx^{*3}z_1^2(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{rx^{*3}z_1^3(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\ &\quad - \frac{cx^{*}y^{*}z_1^2(t)}{1+x^{*2}[1+z_1(t)]^2} - \frac{\beta y^{*}z_2^2(t)}{x^{*}[1+z_1(t)]} \\ &\quad - \frac{cx^{*}y^{*}z_1^2(t)z_2(t)}{1+x^{*2}[1+z_1(t)]^2} \\ &\quad + \frac{\delta\{1+x^{*2}[1+z_1(t)]^2\} - cx^{*}y^{*}[1+z_1(t)]}{\{1+x^{*2}[1+z_1(t)]^2\}[1+z_1(t)]}z_1(t)z_2(t) \end{aligned} \quad (2.62)$$

If $\delta(1+m^2) - cy^{*}M > 0$, and by Lemma 2.3, there exists a $T^{*} > 0$ such that $m \leq x^{*}[1+z_1(t)] \leq M$ and $l \leq y^{*}[1+z_2(t)] \leq L$ for $t > T^{*}$. Then (2.62) implies that

$$\begin{aligned}
\dot{V}_1(z(t)) &\leq \frac{2rx^{*2}z_1^2(t)}{1+x^{*2}[1+z_1(t)]^2} - \frac{2rx^{*3}z_1^2(t)}{K\{1+x_1^{*2}[1+z_1(t)]^2\}} - \frac{cx^*y^*z_1^2(t)}{1+x^{*2}[1+z_1(t)]^2} \\
&\quad + \frac{rx^{*2}z_1^2(t)|z_1(t)|}{1+x^{*2}[1+z_1(t)]^2} + \frac{rx^{*3}z_1^2(t)|z_1(t)|}{K\{1+x_1^{*2}[1+z_1(t)]^2\}} + \frac{2rx^{*3}z_1^2(t)|z_1(t-\tau)|}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\
&\quad + \frac{cx^*y^*z_1^2(t)|z_2(t)|}{1+x^{*2}[1+z_1(t)]^2} + \frac{rx^{*3}z_1^2(t)|z_1(t)||z_1(t-\tau)|}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{\beta y^*z_2^2(t)}{x^*[1+z_1(t)]} \\
&\quad + \frac{\delta\{1+x^{*2}[1+z_1(t)]^2\} - cx^*y^*[1+z_1(t)]}{\{1+x^{*2}[1+z_1(t)]^2\}[1+z_1(t)]}|z_1(t)||z_2(t)| \\
&\quad - \frac{rx^*z_1(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} - \frac{rx^{*3}z_1(t)z_1(t-\tau)}{K\{1+x^{*2}[1+z_1(t)]^2\}} \\
&\leq \frac{2rx^{*2}}{1+m^2}z_1^2(t) - \frac{2rx^{*3}}{K(1+M^2)}z_1^2(t) - \frac{cx^*y^*}{1+M^2}z_1^2(t) + \frac{rx^{*2}}{1+m^2}z_1^2(t)|z_1(t)| \\
&\quad + \frac{rx^{*3}}{K(1+m^2)}z_1^2(t)|z_1(t)| + \frac{2rx^{*3}}{K(1+m^2)}z_1^2(t)|z_1(t-\tau)| \\
&\quad + \frac{cx^*y^*}{1+m^2}z_1^2(t)|z_2(t)| + \frac{rx^{*3}}{K(1+m^2)}z_1(t)^2|z_1(t)||z_1(t-\tau)| - \frac{\beta y^*}{M}z_2^2(t) \\
&\quad + \frac{\delta x^*(1+M^2)}{2m(1+m^2)}[z_1^2(t)+z_2^2(t)] - \frac{cx^*y^*}{2(1+m^2)}[z_1^2(t)+z_2^2(t)] \\
&\quad - \frac{rx^*(1+x^{*2})}{K\{1+x^{*2}[1+z_1(t)]^2\}}z_1(t)\left[z_1(t)-\int_{t-\tau}^t\dot{z}_1(s)ds\right] \\
&\leq \frac{2rx^{*2}}{1+m^2}z_1^2(t) - \frac{2rx^{*3}}{K(1+M^2)}z_1^2(t) - \frac{cx^*y^*}{1+M^2}z_1^2(t) + \frac{rx^*(M+x^*)}{1+m^2}z_1^2(t) \\
&\quad + \frac{rx^{*2}(M+x^*)}{K(1+m^2)}z_1^2(t) + \frac{cx^*(L+y^*)}{1+m^2}z_1^2(t) - \frac{\beta y^*}{M}z_2^2(t) + \frac{\delta x^*(1+M^2)}{2m(1+m^2)}z_1^2(t) \\
&\quad - \frac{cx^*y^*}{2(1+m^2)}z_1^2(t) + \frac{\delta x^*(1+M^2)}{2m(1+m^2)}z_2^2(t) - \frac{cx^*y^*}{2(1+m^2)}z_2^2(t) \\
&\quad - \frac{rx^*(1+x^{*2})}{K(1+M^2)}z_1^2(t) + \frac{2rx^{*3}}{K(1+m^2)}z_1^2(t)|z_1(t-\tau)| \\
&\quad + \frac{rx^{*3}}{K(1+m^2)}z_1(t)^2|z_1(t)||z_1(t-\tau)| \\
&\quad + \frac{rx^*(1+x^{*2})}{K\{1+x^{*2}[1+z_1(t)]^2\}}\int_{t-\tau}^t[1+z_1(s)] \\
&\quad \times \left\{ \frac{2rx^{*2}z_1(t)z_1(s)}{1+x^{*2}[1+z_1(s)]^2} + \frac{rx^{*2}z_1(t)z_1^2(s)}{1+x^{*2}[1+z_1(s)]^2} - \frac{2rx^{*3}z_1(t)z_1(s)}{K\{1+x^{*2}[1+z_1(s)]^2\}} \right. \\
&\quad \left. - \frac{rx^{*3}z_1(t)z_1^2(s)}{K\{1+x^{*2}[1+z_1(s)]^2\}} - \frac{rx^*z_1(t)z_1(s-\tau)}{K\{1+x^{*2}[1+z_1(s)]^2\}} \right. \\
&\quad \left. - \frac{rx^{*3}z_1(t)z_1(s-\tau)}{K\{1+x^{*2}[1+z_1(s)]^2\}} - \frac{2rx^{*3}z_1(t)z_1(s)z_1(s-\tau)}{K\{1+x^{*2}[1+z_1(s)]^2\}} \right\}
\end{aligned}$$

$$\begin{aligned}
& - \frac{rx^{*3}z_1(t)z_1^2(s)z_1(s-\tau)}{K \left\{ 1 + x^{*2} [1 + z_1(s)]^2 \right\}} - \frac{cx^*y^*z_1(t)z_1(s)}{1 + x^{*2} [1 + z_1(s)]^2} \\
& - \frac{cx^*y^*z_1(t)z_2(s)}{1 + x^{*2} [1 + z_1(s)]^2} - \frac{cx^*y^*z_1(t)z_1(s)z_2(s)}{1 + x^{*2} [1 + z_1(s)]^2} \Big\} ds
\end{aligned} \tag{2.63}$$

Then for $t \geq T^* + \tau \equiv \hat{T}$, we have from (2.63) that

$$\begin{aligned}
\dot{V}_1(z(t)) &\leq - \frac{cx^*y^*}{1 + M^2} z_1^2(t) + \frac{rx^*(M + 3x^*)}{1 + m^2} z_1^2(t) + \frac{3rx^{*2}(M + x^*)}{K(1 + m^2)} z_1^2(t) \\
&+ \frac{cx^*(L + y^*)}{1 + m^2} z_1^2(t) + \frac{rx^*(M + x^*)^2}{K(1 + m^2)} z_1^2(t) - \frac{\beta y^*}{M} z_2^2(t) \\
&+ \frac{\delta x^*(1 + M^2)}{2m(1 + m^2)} z_1^2(t) - \frac{cx^*y^*}{2(1 + m^2)} z_1^2(t) - \frac{rx^*(1 + 3x^{*2})}{K(1 + M^2)} z_1^2(t) \\
&+ \frac{\delta x^*(1 + M^2)}{2m(1 + m^2)} z_2^2(t) - \frac{cx^*y^*}{2(1 + m^2)} z_2^2(t) \\
&+ \frac{Mr(1 + x^{*2})}{K(1 + m^2)^2} \int_{t-\tau}^t \left(2rx^{*2} |z_1(t)| |z_1(s)| + \frac{2rx^{*3}}{K} |z_1(t)| |z_1(s)| \right. \\
&+ \frac{rx^*}{K} |z_1(t)| |z_1(s-\tau)| + cx^*y^* |z_1(t)| |z_2(s)| + cx^*y^* |z_1(t)| |z_1(s)| \\
&\quad \left. + \frac{rx^{*3}}{K} |z_1(t)| |z_1(s-\tau)| \right) ds r \\
&+ \frac{rx^*(1 + x^{*2})}{K \left\{ 1 + x^{*2} [1 + z_1(t)]^2 \right\}} \int_{t-\tau}^t [1 + z_1(s)] \\
&\times \left\{ \left[\frac{rx^{*2}z_1(t)z_1(s)}{1 + x^{*2} [1 + z_1(s)]^2} - \frac{rx^{*3}z_1(t)z_1(s)}{K \left\{ 1 + x^{*2} [1 + z_1(s)]^2 \right\}} \right. \right. \\
&- \frac{2rx^{*3}z_1(t)z_1(s-\tau)}{K \left\{ 1 + x^{*2} [1 + z_1(s)]^2 \right\}} - \frac{cx^*y^*z_1(t)z_2(s)}{1 + x^{*2} [1 + z_1(s)]^2} \Big] [1 + z_1(s)] \\
&- \left[\frac{rx^{*2}z_1(t)z_1(s)}{1 + x^{*2} [1 + z_1(s)]^2} - \frac{rx^{*3}z_1(t)z_1(s)}{K \left\{ 1 + x^{*2} [1 + z_1(s)]^2 \right\}} \right. \\
&\quad \left. \left. - \frac{2rx^{*3}z_1(t)z_1(s-\tau)}{K \left\{ 1 + x^{*2} [1 + z_1(s)]^2 \right\}} - \frac{cx^*y^*z_1(t)z_2(s)}{1 + x^{*2} [1 + z_1(s)]^2} \right] \right\} ds
\end{aligned}$$

$$\begin{aligned}
& -\frac{rx^*(1+x^{*2})}{K\left\{1+x^{*2}[1+z_1(t)]^2\right\}} \int_{t-\tau}^t [1+z_1(s)] \\
& \times \left\{ \frac{rx^{*3}z_1(t)z_1(s-\tau)}{K\left\{1+x^{*2}[1+z_1(s)]^2\right\}} [1+z_1(s)]^2 \right. \\
& \quad \left. -\frac{rx^{*3}z_1(t)z_1(s-\tau)}{K\left\{1+x^{*2}[1+z_1(s)]^2\right\}} -\frac{2rx^{*3}z_1(t)z_1(s)z_1(s-\tau)}{K\left\{1+x^{*2}[1+z_1(s)]^2\right\}} \right\} ds \\
\leq & -\frac{cx^*y^*}{1+M^2}z_1^2(t) + \frac{rx^*(M+3x^*)}{1+m^2}z_1^2(t) + \frac{3rx^{*2}(M+x^*)}{K(1+m^2)}z_1^2(t) \\
& + \frac{cx^*(L+y^*)}{1+m^2}z_1^2(t) + \frac{rx^*(M+x^*)^2}{K(1+m^2)}z_1^2(t) - \frac{\beta y^*}{M}z_2^2(t) \\
& + \frac{\delta x^*(1+M^2)}{2m(1+m^2)}z_1^2(t) - \frac{cx^*y^*}{2(1+m^2)}z_1^2(t) - \frac{rx^*(1+3x^{*2})}{K(1+M^2)}z_1^2(t) \\
& + \frac{\delta x^*(1+M^2)}{2m(1+m^2)}z_2^2(t) - \frac{cx^*y^*}{2(1+m^2)}z_2^2(t) \\
& + \frac{Mr(1+x^{*2})}{K(1+m^2)^2} \left[rx^{*2}\tau z_1^2(t) + rx^{*2} \int_{t-\tau}^t z_1^2(s)ds + \frac{rx^{*3}\tau}{K}z_1^2(t) \right. \\
& + \frac{rx^{*3}}{K} \int_{t-\tau}^t z_1^2(s)ds + \frac{rx^*\tau}{2K}z_1^2(t) + \frac{rx^*}{2K} \int_{t-\tau}^t z_1^2(s-\tau)ds + \frac{rx^{*3}\tau}{2K}z_1^2(t) \\
& + \frac{rx^{*3}}{2K} \int_{t-\tau}^t z_1^2(s-\tau)ds + \frac{cx^*y^*\tau}{2}z_1^2(t) + \frac{cx^*y^*}{2} \int_{t-\tau}^t z_1^2(s)ds \\
& + \frac{cx^*y^*\tau}{2}z_1^2(t) + \frac{cx^*y^*}{2} \int_{t-\tau}^t z_2^2(s)ds \left. \right] \\
& + \frac{M^2r(1+x^{*2})}{Kx^*(1+m^2)^2} \int_{t-\tau}^t \left[rx^{*2}|z_1(t)||z_1(s)| + \frac{rx^{*3}}{K}|z_1(t)||z_1(s)| \right. \\
& \quad \left. + \frac{2rx^{*3}}{K}|z_1(t)||z_1(s-\tau)| + cx^*y^*|z_1(t)||z_2(s)| \right] ds \\
& + \frac{Mr(1+x^{*2})}{K(1+m^2)^2} \int_{t-\tau}^t \left[rx^{*2}|z_1(t)||z_1(s)| + \frac{rx^{*3}}{K}|z_1(t)||z_1(s)| \right. \\
& \quad \left. + \frac{2rx^{*3}}{K}|z_1(t)||z_1(s-\tau)| + cx^*y^*|z_1(t)||z_2(s)| \right] ds \\
& + \frac{M^3r^2x^*(1+x^{*2})}{K^2(1+m^2)^2} \int_{t-\tau}^t |z_1(t)||z_1(s-\tau)| ds \\
& + \frac{Mr^2x^{*3}(1+x^{*2})}{K^2(1+m^2)^2} \int_{t-\tau}^t |z_1(t)||z_1(s-\tau)| ds \\
& + \frac{2Mr^2x^{*3}(1+x^{*2})}{K^2(1+m^2)^2} \int_{t-\tau}^t \{|z_1(t)||z_1(s-\tau)|[1+z_1(s)] \\
& \quad + |z_1(t)||z_1(s-\tau)|\} ds
\end{aligned}$$

$$\begin{aligned}
&\leq - \left[\frac{rx^*(1+3x^{*2})}{K(1+M^2)} + \frac{cx^*y^*}{2(1+m^2)} + \frac{cx^*y^*}{1+M^2} - \frac{rx^*(M+3x^*)}{1+m^2} \right. \\
&\quad - \frac{3rx^{*2}(M+x^*)}{K(1+m^2)} - \frac{rx^*(M+x^*)^2}{K(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} - \frac{cx^*(L+y^*)}{1+m^2} \\
&\quad - \frac{Mr(1+x^{*2})\tau}{K(1+m^2)^2} \left(\frac{3rx^{*2}}{2} + \frac{9rx^{*3}}{2K} + \frac{rx^*}{2K} + \frac{3cx^*y^*}{2} \right) \\
&\quad \left. - \frac{M^2r(1+x^{*2})\tau}{Kx^*(1+m^2)^2} \left(\frac{rx^{*2}}{2} + \frac{5rx^{*3}}{2K} + \frac{cx^*y^*}{2} \right) - \frac{M^3r^2x^*(1+x^{*2})\tau}{2K^2(1+m^2)^2} \right] z_1^2(t) \\
&\quad - \left[\frac{\beta y^*}{M} + \frac{cx^*y^*}{2(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} \right] z_2^2(t) \\
&\quad + \frac{Mr(1+x^{*2})}{K(1+m^2)^2} \left[\left(\frac{3rx^{*2}}{2} + \frac{3rx^{*3}}{2K} + \frac{cx^*y^*}{2} \right) \int_{t-\tau}^t z_1^2(s) ds \right. \\
&\quad \left. + cx^*y^* \int_{t-\tau}^t z_2^2(s) ds + \left(\frac{rx^*}{2K} + \frac{3rx^{*3}}{K} \right) \int_{t-\tau}^t z_1^2(s-\tau) ds \right] \\
&\quad + \frac{M^2r(1+x^{*2})}{Kx^*(1+m^2)^2} \left[\left(\frac{rx^{*2}}{2} + \frac{rx^{*3}}{2K} \right) \int_{t-\tau}^t z_1^2(s) ds \right. \\
&\quad \left. + \frac{cx^*y^*}{2} \int_{t-\tau}^t z_2^2(s) ds + \frac{2rx^{*3}}{K} \int_{t-\tau}^t z_1^2(s-\tau) ds \right] \\
&\quad + \frac{M^3r^2x^*(1+x^{*2})}{2K^2(1+m^2)^2} \int_{t-\tau}^t z_1^2(s-\tau) ds \tag{2.64}
\end{aligned}$$

Let

$$\begin{aligned}
V_2(z(t)) &= \frac{Mrx^*(1+x^{*2})(3Krx^*+3rx^{*2}+Kcy^*+MrK+Mrx^*)}{2K^2(1+m^2)^2} \int_{t-\tau}^t \int_s^t z_1^2(\gamma) d\gamma ds \\
&\quad + \frac{Mrcy^*(1+x^{*2})(2x^*+M)}{2K(1+m^2)^2} \int_{t-\tau}^t \int_s^t z_2^2(\gamma) d\gamma ds \\
&\quad + \frac{Mr^2x^*(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2} \int_{t-\tau}^t \int_s^t z_1^2(\gamma-\tau) d\gamma ds \tag{2.65}
\end{aligned}$$

then

$$\begin{aligned}
\dot{V}_2(z(t)) &= \frac{Mrx^*\tau(1+x^{*2})(3Krx^*+3rx^{*2}+Kcy^*+MrK+Mrx^*)}{2K^2(1+m^2)^2}z_1^2(t) \\
&\quad - \frac{Mrx^*(1+x^{*2})(3Krx^*+3rx^{*2}+Kcy^*+MrK+Mrx^*)}{2K^2(1+m^2)^2} \int_{t-\tau}^t z_1^2(s)ds \\
&\quad + \frac{Mrcy^*\tau(1+x^{*2})(2x^*+M)}{2K(1+m^2)^2}z_2^2(t) \\
&\quad - \frac{Mrcy^*(1+x^{*2})(2x^*+M)}{2K(1+m^2)^2} \int_{t-\tau}^t z_2^2(s)ds \\
&\quad + \frac{Mr^2x^*\tau(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2}z_1^2(t-\tau) \\
&\quad - \frac{Mr^2x^*(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2} \int_{t-\tau}^t z_1^2(s-\tau)ds
\end{aligned} \tag{2.66}$$

and then we have from (2.64) and (2.66) that for $t \geq \hat{T}$

$$\begin{aligned}
\dot{V}_1(z(t)) + \dot{V}_2(z(t)) &\leq - \left[\frac{rx^*(1+3x^{*2})}{K(1+M^2)} + \frac{cx^*y^*}{2(1+m^2)} + \frac{cx^*y^*}{1+M^2} \right. \\
&\quad - \frac{rx^*(M+3x^*)}{1+m^2} - \frac{3rx^{*2}(M+x^*)}{K(1+m^2)} - \frac{rx^*(M+x^*)^2}{K(1+m^2)} \\
&\quad - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} - \frac{cx^*(L+y^*)}{1+m^2} \\
&\quad \left. - \frac{Mrx^*\tau(1+x^{*2})(3Krx^*+3rx^{*2}+Kcy^*+MrK+Mrx^*)}{2K^2(1+m^2)^2} \right. \\
&\quad - \frac{Mr(1+x^{*2})\tau}{K(1+m^2)^2} \left(\frac{3rx^{*2}}{2} + \frac{9rx^{*3}}{2K} + \frac{rx^*}{2K} + \frac{3cx^*y^*}{2} \right) \\
&\quad - \frac{M^2r(1+x^{*2})\tau}{Kx^*(1+m^2)^2} \left(\frac{rx^{*2}}{2} + \frac{5rx^{*3}}{2K} + \frac{cx^*y^*}{2} \right) \\
&\quad \left. - \frac{M^3r^2x^*(1+x^{*2})\tau}{2K^2(1+m^2)^2} \right] z_1^2(t)
\end{aligned}$$

$$\begin{aligned}
& - \left[\frac{\beta y^*}{M} + \frac{cx^*y^*}{2(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} \right. \\
& \quad \left. - \frac{Mrcy^*\tau(1+x^{*2})(2x^*+M)}{2K(1+m^2)^2} \right] z_2^2(t) \\
& + \frac{Mr^2x^*\tau(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2} z_1^2(t-\tau)
\end{aligned} \tag{2.67}$$

Let

$$V_3(z(t)) = \frac{Mr^2x^*\tau(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2} \int_{t-\tau}^t z_1^2(s) ds \tag{2.68}$$

then

$$\begin{aligned}
\dot{V}_3(z(t)) &= \frac{Mr^2x^*\tau(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2} z_1^2(t) \\
&\quad - \frac{Mr^2x^*\tau(1+x^{*2})(1+6x^{*2}+4Mx^*+M^2)}{2K^2(1+m^2)^2} z_1^2(t-\tau)
\end{aligned} \tag{2.69}$$

Now define a Lyapunov functional $V(z(t))$ as

$$V(z(t)) = V_1(z(t)) + V_2(z(t)) + V_3(z(t)) \tag{2.70}$$

then we have from (2.67) and (2.69) that for $t \geq \hat{T}$

$$\begin{aligned}
\dot{V}(z(t)) &\leq - \left[\frac{rx^*(1+3x^{*2})}{K(1+M^2)} + \frac{cx^*y^*}{1+M^2} + \frac{cx^*y^*}{2(1+m^2)} - \frac{rx^*(M+3x^*)}{1+m^2} \right. \\
&\quad - \frac{3rx^{*2}(M+x^*)}{K(1+m^2)} - \frac{rx^*(M+x^*)^2}{K(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} - \frac{cx^*(L+y^*)}{1+m^2} \\
&\quad \left. - \frac{Mr^2x^*\tau(1+x^{*2})(3Kx^*+9x^{*2}+MK+5Mx^*+1+M^2)}{K^2(1+m^2)^2} \right. \\
&\quad \left. - \frac{Mrcy^*\tau(1+x^{*2})(4x^*+M)}{2K(1+m^2)^2} \right] z_1^2(t) \\
&\quad - \left[\frac{\beta y^*}{M} + \frac{cx^*y^*}{2(1+m^2)} - \frac{\delta x^*(1+M^2)}{2m(1+m^2)} - \frac{Mrcy^*\tau(1+x^{*2})(2x^*+M)}{2K(1+m^2)^2} \right] z_2^2(t) \\
&\equiv -\zeta z_1^2(t) - \eta z_2^2(t)
\end{aligned} \tag{2.71}$$

Then it follows from (2.57) and (2.58) that $\zeta > 0$ and $\eta > 0$. Let $w(s) = \hat{N}s^2$ where $\hat{N} = \min\{\zeta, \eta\}$, then w is nonnegative continuous on $[0, \infty)$, $w(0) = 0$, and $w(s) > 0$ for $s > 0$. It follows from (2.71) that for $t \geq \hat{T}$

$$\dot{V}(z(t)) \leq -\hat{N}[z_1^2(t) + z_2^2(t)] = -\hat{N}\|z(t)\|^2 = -w(\|z(t)\|) \quad (2.72)$$

Now, we want to find a function u such that $V(z(t)) \geq u(\|z(t)\|)$. It follows from (2.61), (2.65), and (2.68) that

$$V(z(t)) \geq \{z_1(t) - \ln[1 + z_1(t)]\} + \{z_2(t) - \ln[1 + z_2(t)]\} \quad (2.73)$$

then by Theorem 2.1, we have that $u(s) = \tilde{N}s^2$ and $V(z(t)) \geq u(\|z(t)\|)$. So the equilibrium point E^* of the system (2.1) is globally asymptotically stable with $p(x) = \frac{cx^2}{1+x^2}$

Remark 2.1 By Theorem 2.1, Theorem 2.2, and Theorem 2.3 we can assume that

$$\begin{aligned} V(z(t)) &= \{z_1(t) - \ln[1 + z_1(t)]\} + \{z_2(t) - \ln[1 + z_2(t)]\} \\ &\quad + a(p(x)) \frac{Mrx^*}{2K^2} \int_{t-\tau}^t \int_s^t z_1^2(\gamma) d\gamma ds \\ &\quad + b(p(x)) \frac{Mr^2x^*}{2K^2} \left[\int_{t-\tau}^t \int_s^t z_1^2(\gamma - \tau) d\gamma ds + \tau \int_{t-\tau}^t z_1^2(s) ds \right] \\ &\quad + c(p(x)) \frac{Mrcy^*}{2K} \int_{t-\tau}^t \int_s^t z_2^2(\gamma) d\gamma ds \end{aligned}$$

Then

(i) for $p(x) = cx$ in the Holling-Type I model

$$a(p(x)) = 0, \quad b(p(x)) = 1, \quad c(p(x)) = 1$$

(ii) for $p(x) = \frac{cx}{1+x}$ in the Holling-Type II model

$$\begin{aligned} a(p(x)) &= \frac{r(1+x^*)}{(1+m)^2} \\ b(p(x)) &= \frac{(1+x^*)(M+1+2x^*)}{(1+m)^2} \\ c(p(x)) &= \frac{1+x^*}{(1+m)^2} \end{aligned}$$

(iii) for $p(x) = \frac{cx^2}{1+x^2}$ in the Holling-Type III model

$$\begin{aligned} a(p(x)) &= \frac{(1+x^{*2})(3Krx^* + 3rx^{*2} + Kcy^* + MKr + Mrx^*)}{(1+m^2)^2} \\ b(p(x)) &= \frac{(1+x^{*2})(1+6x^{*2} + 4Mx^* + M^2)}{(1+m^2)^2} \\ c(p(x)) &= \frac{(1+x^{*2})(2x^* + M)}{(1+m^2)^2} \end{aligned}$$

3 Examples

Example 3.1 In the Holling-Type I we consider the system

$$\begin{aligned} \dot{x}(t) &= x(t)[3 - 10x(t-\tau) - 15y(t)] \\ \dot{y}(t) &= y(t)\left[1 - 6\frac{y(t)}{x(t)}\right] \end{aligned} \tag{3.1}$$

where $r = 3$, $K = \frac{3}{10}$, $c = 15$, $\delta = 1$, $\beta = 6$, and $E^* = (\frac{6}{25}, \frac{1}{25})$. Then

$$\begin{aligned} r - cL &= 2.24246 > 0 \\ \delta x^* - cy^*M &= 0.0582 > 0 \\ \frac{rx^*}{K} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mr^2x^*\tau}{K^2} - \frac{Mrcy^*\tau}{2K} &= 1.5940 > 0 \\ \frac{\beta y^*}{M} + \frac{cy^*}{2} - \frac{\delta x^*}{2m} - \frac{Mrcy^*\tau}{2K} &= 0.0103 > 0 \end{aligned}$$

whenever $\tau = \frac{1}{300}$. Consequently, by Theorem 2.1, we conclude that the unique positive equilibrium point E^* of the system (3.1) is globally asymptotically stable. The trajectory of the system (3.1) is depicted in Figure 3.1.




Figure 3.1: The trajectory of the system (3.1) with $\tau = \frac{1}{300}$.

Example 3.2 In the Holling-Type II we consider the system

$$\begin{aligned}\dot{x}(t) &= x(t) \left[3 - 10x(t-\tau) - 15 \frac{y(t)}{1+x(t)} \right] \\ \dot{y}(t) &= y(t) \left[1 - 6 \frac{y(t)}{x(t)} \right]\end{aligned}\tag{3.2}$$

where $r = 3$, $K = \frac{3}{10}$, $c = 15$, $\delta = 1$, $\beta = 6$, and $E^* = (\frac{1}{4}, \frac{1}{24})$. Then

$$\begin{aligned}r - cL &= 2.24246 > 0 \\ \delta x^* (1+m) - cy^* M &= 0.08843 > 0 \\ \frac{rx^*(1+2x^*)}{K(1+M)} + \frac{cy^*}{2(1+m)} - \frac{rx^*}{1+m} - \frac{\delta x^*(1+M)}{2m(1+m)} - \frac{rx^*(x^*+M)}{K(1+m)} \\ - \frac{Mr^2x^*\tau(1+x^*)(K+3x^*+1+M)}{K^2(1+m)^2} - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} &= 0.05316 > 0 \\ \frac{\beta y^*}{M} + \frac{cy^*}{2(1+m)} - \frac{\delta x^*(1+M)}{2m(1+m)} - \frac{Mrcy^*\tau(1+x^*)}{2K(1+m)^2} &= 0.00475 > 0\end{aligned}$$

whenever $\tau = \frac{1}{300}$. Consequently, by Theorem 2.2, we conclude that the unique positive equilibrium point E^* of the system (3.2) is globally asymptotically stable. The trajectory of the system (3.2) is depicted in Figure 3.2.




Figure 3.2: The trajectory of the system (3.2) with $\tau = \frac{1}{300}$.

4 Conclusion

In this paper, we obtain that the sufficient condition for the global stability of the Leslie-Gower predator-prey system in Holling-Type I, Holling-Type II, and Holling-Type III models with time delay, respectively. But we believe that the global stability of the predator-prey model with time delay with all different functional response of the predator, $p(x)$, for instance, $p(x) = mx$, $p(x) = \frac{mx}{a+x}$, $p(x) = \frac{mx^2}{a+x^2}$, $p(x) = mx^c$, $0 < c \leq 1$, or $p(x) = m(1 - e^{-cx})$ will be an important topic for future study.

References

- [1] He, X. Z., Stability and Delays in a Predator-Prey System, *J. Math. Anal. Appl.* **198**, 355-370 (1996).
- [2] Ho, C. P. and Huang S. T., Global Stability for a Predator-Prey System with Predator Self-Limitation, *Tunghai Science* **3**, 3-27 (2001).

- [3] Ho, C. P. and Jiang J. M., Stabilization of Predator-Prey Systems by Refuge, Thesis, Tunghai University, Taiwan (1997).
- [4] Ho, C. P. and Liang, C. L., Global Stability for the Leslie-Gower Predator-Prey System with Time-Delay, Thesis, Tunghai University, Taiwan (2002).
- [5] Ho, C. P. and Lin K. S., Global Stability for a Class of Predator-Prey Systems with Ratio-Dependence, *Tunghai Science* **3**, 113-133 (2001).
- [6] Hsu, S. B., On Global Stability of a Predator-Prey System, *Math. Biosci.* **39**, 1-10 (1978).
- [7] Hsu, S. B. and Huang T. W., Global Stability for a Class of Predator-Prey System, *SIAMJ. Appl. Math.* **55**, 763-783 (1995).
- [8] Korobeinikov, A., A Lyapunov Function for Leslie-Gower Predator-Prey Models, *Appl. Math. Letters* **14**, 697-699 (2001).
- [9] Kuang, Y., Convergence Results in a Well-Known Delayed Predator-Prey System, *J. Math. Anal. Appl.* **204**, 840-853 (1996).
- [10] Kuang, Y., Delay Differential Equations with Applications in Population Dynamics, Academic Press, Boston (1993).
- [11] Kuang, Y., Global Stability of Gause-Type Predator-Prey Systems, *J. Math. Biol.* **28**, 463-474 (1990).
- [12] Kuang, Y. and Smith, H. L., Global Stability for Infinite Delay Lotka-Volterra Type Systems, *J. Differential Equations* **103**, 221-246 (1993).
- [13] Perko, L., Differential Equations and Dynamical Systems, Springer-Verlag, New York, Inc (1991).
- [14] Wang, W. and Ma, Z, Harmless Delays for Uniform Persistence, *J. Math. Anal. Appl.* **158**, 256-268. (1991).

具時滯參數且函數反應為 Holling 型式的 Leslie-Gower 捕食系統之整體穩定性

蔡翰征* 何肇寶

摘要

本篇討論文主要探討具時滯參數且函數反應為 Holling 型式的 Leslie-Gower 捕食系統之整體穩定性。首先，我們利用 Lyapunov 函數來分別討論不同 Holling 型式時具時滯參數的 Leslie-Gower 捕食系統之整體穩定性。最後，我們舉例說明我們的結果。

關鍵詞：Leslie-Gower 捕食系統，整體穩定性，時滯參數，Holling 型式，Lyapunov 函數。

* 東海大學數學系

具時滯參數且函數反應為 Holling 型式的 Leslie-Gower 捕食系統之整體穩定性

蔡翰征* 何肇寶

摘要

本篇討論文主要探討具時滯參數且函數反應為 Holling 型式的 Leslie-Gower 捕食系統之整體穩定性。首先，我們利用 Lyapunov 函數來分別討論不同 Holling 型式時具時滯參數的 Leslie-Gower 捕食系統之整體穩定性。最後，我們舉例說明我們的結果。

關鍵詞：Leslie-Gower 捕食系統，整體穩定性，時滯參數，Holling 型式，Lyapunov 函數。

* 東海大學數學系