

陸、参考文献

- 江文德。1998。簡介大豆中的異黃酮素。食品工業，30(9):6-12。
- 青山敏明。1999。大豆たん白質および大豆ペプチドの生理機能研究。食品と開発，34(7):15-17。
- Adlercreutz, H., Mousavi, Y., Clark, J., Höckerstedt, K., Hämäläinen, E., Wähälä, K., Mäkelä, T. and Hase, T. (1992). Dietary phytoestrogens and cancer: in vitro and in vivo studies. *Journal of Steroid Biochemistry and Molecular Biology*, 41:331-337.
- Adlercreutz, H., Fotsis, T., Lampe, J., Wähälä, K., Mäkelä, T., Brunow, G. and Hase, T. (1993a). Quantitative determination of lignans and isoflavonoids in plasma of omnivorous and vegetarian women by isotope dilution gas-chromatography mass-spectrometry. *Scandinavian Journal of Clinical and Laboratory Investigation*, 53:5-18.
- Adlercreutz, H., Bannwart, C., Wähälä, K., Mäkelä, T., Brunow, G., Hase, T., Arosemena, P.J., Kellis, J.T. and Vickery, L.E. (1993b). Inhibition of human aromatase by mammalian lignans and isoflavonoid phytoestrogens. *Journal of Steroid Biochemistry and Molecular Biology*, 44:127-153.
- Adlercreutz, H., Fotsis, T., Kurzer, M.S., Wähälä, K., Mäkelä, T. and Hase, T. (1995). Isotope dilution gas chromatographic-mass spectrometric method for the determination of unconjugated lignans and isoflavonoids in human feces, with preliminary results in omnivorous and vegetarian women. *Analytical Biochemistry*, 225:101-108.
- Amarowicz, R., Wanasundara, U., Wanasundara, J. and Shahidi, F. (1993). Antioxidant activity of ethanolic extracts of flaxseed in a β -carotene-linoleate model system. *Journal of Food Lipids*, 1:111-117.
- Arditi, T., Meredith, T. (1999). Soy isoflavones and saponins: a natural soy solution to age-old problems. *Agro-food-Industry Hi-Tech*-July/August pp12-14
- Axelsson, M. and Setchell, K.D.R. (1980). Conjugation of lignans in human urine. *FEBS Letters*, 122:49-53.
- Axelsson, M. and Setchell, K.D.R. (1981). The excretion of lignans in rats – evidence for an intestinal bacterial source for this new group of compounds. *FEBS Letters*, 123:337-342.
- Axelsson, M., Sjövall, J., Gustafsson, B.E. and Setchell, K.D.R. (1982a). Origin of lignans in mammals and identification of a precursor from plants. *Nature*, 298:659-660.

- Axelsson, M., Kirk, D.N., Farrant, R.D., Cooley, G., Lawson, A.M. and Seteshell, K.D.R. (1982b). The identification of the weak oestrogen equol [7-hydroxy-3-4'-hydroxyphenylchroman] in human urine. *Biochemical Journal*, 201:353-357.
- Axelsson, M., Sjövall, J., Gustafsson, B.E. and Setechell, K.D.R. (1984). Soya – a dietary source of the nonsteroidal oestrogen equol in man and animals. *Journal of Endocrinology*, 102:49-56.
- Ayres, D.C. and Loike, J.D. (1990). *Lignans: Chemical, biological, and clinical properties*, Cambridge University Press, Cambridge, pp. 1-11; 85-112.
- Bakke, J.E. and Klosterman, H.J. (1956). A new diglucoside from flaxseed. *Proceedings of the North Dakota Academy of Science*, 10:18-22.
- Bambagiotti-Alberti, M., Coran, S.A., Ghiara, C., Giannellini, V. And Raffaelli, A. (1994). Revealing the mammalian lignan precursor secoisolariciresinol diglucoside in flaxseed by ion spray mass-spectrometry. *Rapid Communications in Mass Spectrometry*, 8:595-598.
- Begum, A.N., Nivolle, C., Mila, I., Lapierre, C., Nagano, K., Fulushima, K., Heinonen, S.M., Adlercreutz, H., Rémésy, C. and Scalbert, A. (2003). Dietary lignans are precursors of mammalian lignans in rats. *Journal of nutrition*, 134:120-127.
- BeMiller, J.N. (1973). Quince seed, psyllium seed, flaxseed and okra gums. In: R.L. Whistler and J.N. BeMiller (Eds.), *Industrial gums*. Academic Press, New York, pp. 331-337.
- Bhatty, R.S. (1997). Nutrient composition of whole flaxseed and flaxseed meal. In: S.C. Cunnane and L.U. Thompson (Eds.), *Flaxseed in human nutrition*. AOCS Press, Champaign, IL, pp. 22-42.
- Borriello, S.P., Setchell, K.D.R., Axelsson, M. and Lawson, A.M. (1985). Production and metabolism of lignans by the human faecal flora. *Journal of applied bacteriology*, 58:37-43.
- Bowey, E., Adlercreutz, H. and Rowland, I. (2003). Metabolism of isoflavone and lignans by the gut microflora: a study in germ-free and human flora associated rats. *Food and Chemical Toxicology*, 41:631-636.
- Bradford, M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72:248-254.

- Carmela, G., Federica, B., Angelo, S. and Giuseppe, Z. (2001). Purification and characterisation of a b-glucosidase abundantly expressed in ripe sweet cherry (*Prunus abium* L.) fruit. *Plant Science* 160: 795–805
- Charlet, S., Bensaddek, L., Raynaud, S., Gillet F., Mesnard, F. and Fliniaux, M.A. (2002). An HPLC method for the quantification of anhydrosecoisolariciresinol. Application to the evaluation of flax lignan content. *Plant physiology and biochemistry*, 40:225-229.
- Chen, J., Hui, E., Ip, T. and Thompson, L.U. (2004). Dietary flaxseed enhances the inhibitory effect of tamoxifen on the growth of estrogen-dependent human breast cancer (MCF-7) in nude mice. *Clinical Cancer Research*, 10:7703-7711.
- Chen, H.H., Xu, S.Y. and Wang, Z. (2006). Gelation properties of flaxseed gum. *Journal of Food Engineering*, 77:295-303.
- Chen, H.M., Muramoto, K. and Yamauchi, F. (1995). Structural analysis of antioxidative peptides from soybean β -conglycinin. *J. Agric. Food Chem.*, 43: 574-578
- Christina Strandås, Afaf Kamal-Eldina, Roger Andersson and Per Åmana. (2008). Composition and properties of flaxseed phenolic oligomers. *Food Chem.* Volume 110, Issue 1, 1 September 2008, Pages 106-112 .
- Chung, M.W.Y., Lei, B. and Li-Chan, E.C.Y. (2005). Isolation and structural characterization of the major protein fraction from NorMan flaxseed (*Linum usitatissimum* L.). *Food Chemistry*, 90:271-279.
- Clavel, T., Henderson, G., Alpert, C.A, Philippe, C., Rigottier-Gois, L., Doré, J. and Blaut, M. (2005). Intestinal bacterial communities that produce active estrogen-like compounds enterodiol and enterolactone in humans. *Appl Environ Microbiol*, 71:6077–6085
- Clavel, T., Henderson, G., Engst, W., Doré, J. and Blaut, M. (2006a) Phylogeny of human intestinal bacteria that activate the dietary lignan secoisolariciresinol diglucoside. *FEMS Microbiol*, 55:471–478
- Clavel, T., Borrmann, D., Braune, A., Doré, J. and Blaut, M. (2006b) Occurrence and activity of human intestinal bacteria involved in the conversion of dietary lignans. *Anaerobe*, 12:140–147.
- Coskuner, Y. and Karababa, E. (2007). Some physical properties of flaxseed (*Linum usitatissimum* L.). *Journal of Food Engineering*, 78:1067-1073.
- Cui, W., Kenaschuk, E. and Mazza, G. (1996). Influence of genotype on

- chemical composition and rheological properties of flaxseed gums. *Food Hydrocolloids*, 10:221-227.
- Czarnecki, S.K. and Kritchevsky, D. (1992). Dietary protein and atherosclerosis. In: G.U. Liepa, D.C. Bietz, A.C. Beyene, and M.A. Gorman (Eds.), *Dietary proteins: how they alleviate disease and promote better health*. Champaign, American Oil Chemists' Society, pp. 42-56.
- Dabrosan, C., Chen, J., Wang, L. And Thompson, .U. (2002). Flaxseed inhibits metastasis and decreases extracellular vascular endothelial growth factor in human breast cancer xenografts. *Cancer Letter*, 185:31-37.
- Dabrowski, K.J. and Sosulski, F.W. (1984). Composition of free and Hydrolyzable acids in defatted flours of ten oilseeds. *Journal of Agricultural and Food Chemistry*, 32:128-130.
- Daun, J.K. and DeClercq, D.R. (1994). Sixty years of Canadian flaxseed quality suverys at the Grain Research Laboratory. *Proceedings Flax Institute*, 55:192-200.
- Degenhardt, A., Habben, S. and Winterhalter, P. (2002). Isolation of the lignan secoisolariciresinol diglucoside from flaxseed (*Linum usitatissimum L.*) by high-speed counter-current chromatography. *Journal of chromatography A*, 943:299-302.
- Denis, L., Morton, M.S. and Griffiths, K. (1999). Diet and its preventive role in prostatic disease. *European Urology*, 35:377-387.
- Dobbins, T.A. and Wiley, D.B. (2003). Process for recovering secoisolariciresinol diglycoside from de-fatted flaxseed. United States patent 20030216553.
- Eliasson, C., Kamal-Eldin, A., Andersson, R. and Åman, P. (2003). High-performance liquid chromatographic analysis of secoisolariciresinol diglycoside and hydroxycinnamic acid glucosides in flaxseed by alkaline extraction. *Journal of chromatography A*, 1012:151-159.
- Fedeniuk, R.W. and Biliaderis, C.G. (1994). Composition and physicochemical properties of linseed mucilage. *Journal of Agriculture and Food Chemistry*, 42:240-247.
- Freeman, T.P. (1995). Structure of flaxseed. In: S.C. Cunnane and L.U. Thompson (Eds.), *Flaxseed in human nutrition*. AOCS Press, Champaign, IL, pp. 11-21.
- Friedman, M., Brandon, D.L., Bates, A.H. and Hymowitz, T. (1991).

- Comparison of a commercial soybean cultivar and an isolate lacking the Kunitz trypsin inhibitor: composition, nutritional value, and effects of heating. *Journal of Agricultural and Food Chemistry*, 39:327-335.
- Haggans, C.J., Hutchins, A.M., Olson, M.A., Thomas, W., Martini, M.C. and Slavin, J.L. (1999). Effect flaxseed consumption on urinary estrogen metabolites in postmenopausal women. *Nutrition and Cancer*, 33: 188-195.
- Harris, R.K. and Haggerty, W.J. (1993). Assays for potentially anticarcinogenic phytochemicals in Flaxseed. *Cereal foods world*, 38:147-151.
- Heinonen, S., Nurmi, T., Liukkonen, K., Poutanen, K., Wähälä, K., Deyama, T., Nishibi, S., and Adlercreutz, H. (2001). In vitro metabolism of plant lignans: New precursors of mammalian lignans enterolactone and enterodiol. *Journal of Agricultural and Food Chemistry*, 49:3178-3186.
- Heintzman, R. and Bennett, M.L. (2003). Process for extraction and stabilization of phytoestrogens from flaxseed and product therefrom. United States Patent 20030060420.
- Hsiang-Lin, C., Hui-Yu, H. and Cheng-Chun, C. (2006). Transformation of isoflavone phytoestrogens during the fermentation of soymilk with lactic acid bacteria and bifidobacteria. *Food Microbiology* 23 :772–778.
- Hutchins, A.M., Martini, M.C., Olson, B.A., Thomas, W. and Slavin, J.L. (2000). Flaxseed influences urinary lignan excretion in a dose-dependent manner in postmenopausal women. *Cancer Epidemiology Biomarkers and Prevention*, 9:1113-1118.
- James L. S., Carolina A. V. and Leslie P. (2005). Activation of Flaxseed Lignans by Lactic Acid Bacteria and Bifidobacteria. Department of Food Science University of Wisconsin-Madison.
- Jenab, M. and Thompson, L.U. (1996). The influence of flaxseed and lignans on colon carcinogenesis and beta-glucuronidase activity. *Carcinogenesis*, 17:1343-1348.
- Jenkins, D.J., Kendall, C.W., Vidgen, E., Agarwal, S., Rao, A.V., Rpsenberg, R.S., Diamandis, E.P., Novokmet, R., Mehling, C.C., Perera, T., Griffin, L.C. and Cunnane, S.C. (1999). Health aspects of partially defatted flaxseed, including effects on serum lipids, oxidative measures, and ex vivo androgen and progestin activity: a controlled crossover trial. *American Journal of Clinical Nutrition*, 69:395-402.

- Johnsson, P., Kamal-Eldin, A., Lundgren, L.N. and Åman, P. (2000). HPLC method for analysis of secoisolariciresinol diglucoside in flaxseeds. *Journal of Agricultural and Food Chemistry*, 48:5216-5219.
- Johnsson, P., Peerlkamp, N., Kamal-Eldin, A., Andersson, R.E., Andersson, R., Lundgren, L.N. and Åman, P. (2002). Polymeric fractions containing phenol glucosides in flaxseed. *Food Chemistry*, 76:207-212.
- Johnsson, P. (2004). Phenolic compounds in flaxseed: Chromatographic and spectroscopic analyses of glucosidic conjugates. *A PhD. Thesis from Department of Food Science*, Uppsala, Swedish University of Agricultural Sciences, pp. 1-36.
- Kamal-Eldin, Peerlkamp, N., A., Johsson, P., Andersson, R., Andersson, R.E., Lundgren, L.N. and Åman, P. (2001). An oligomer from flaxseed composed of secoisolariciresinol diglucoside and 3-hydroxy-3methyl glutaric acid residues. *Phytochemistry*, 58:587-590.
- Kitts, D.D., Yuan, Y.V., Wijewickreme, A.N. and Thompson, L.U. (1999). Antioxidant activity of the flaxseed lignan secoisolariciresinol diglucoside and its mammalian lignan metabolites enterodiol and enterolactone. *Molecular and Cellular Biochemistry*, 202:91-100.
- Klosterman, H.J. and Smith, F. (1954). The isolation of β -hydroxy- β -methylglutaric acid from the seed of flax (*Linum usitatissimum*). *Journal of the American Chemical Society*, 76:1229-1230.
- Klosterman, H.J., Smith, F. and Claggett, C.O. (1955). The constitution of Linocinnamarin. *Journal of the American Chemical Society*, 77:420-421.
- Kozłowska, H., Zadernowski, R. and Sosulaki, F.E. (1983). Phenolic acid in oilseed flours. *Nahrung*, 27:449-453.
- Kritchevsky, D. (1995). Fibre effects on hyperlipidemia. In: S.C. Cunnane and L.U. Thompson (Eds.), *Flaxseed in Human Nutrition*. AOCS Press, Champaign, IL, pp. 174-186.
- Kuijsten, A., Arts, I.C., Vree, T.B. and Hollman, P.C. (2005). Pharmacokinetics of enterolignans in healthy men and women consuming a single dose of secoisolariciresinol diglucoside. *Journal of Nutrition*, 135:795-801.
- Kuijsten, A., Arts, I.C., Hollman, P.C., van't Vree, T.B. and Kampman, E. (2006). Plasma enterolignans are associated with lower colorectal adenoma risk. *Cancer Epidemiology, Biomarker and Prevention*,

15:1132-1136.

- Lampe, J.W., Martini, M.C., Kurzer, M.S., Adlercreutz, H. and Slavin, J.L. (1994). Urinary lignan and isoflavonoid excretion in premenopausal women consuming flaxseed powder. *American Journal of Clinical Nutrition*, 60:122-128.
- Li, D., Tee, J.A., Thompson, L.U. and Yan, L. (1999). Dietary supplementation with secoisolariciresinol diglucoside (SDG) reduces experimental metastasis of melanoma cells in mice. *Cancer Letters*, 142:91-96.
- Liggins, J., Crimwood, R. and Bingham, S.S. (2000). Extraction and quantification of lignan phytoestrogens in food and human samples. *Analytical Biochemistry*, 287:102-109.
- Lin, X., Switzer, B.R. and Demark-Wahnefried, W. (2001). Effect of mammalian lignans on the growth of prostate cancer cell lines. *Anticancer Research*, 21:3995-4000.
- Lin, X., Gingrich, J.R., Bao, W., Li, J., Haroon, Z.A. and Demark-Wahnefried, W. (2002). Effect of flaxseed supplementation on prostatic carcinoma in transgenic mice. *Urology*, 60:919-924.
- Lupien, P.J., Moorjani, S., Brun, D. and Biemann, I. (1979). Effects of 3-hydroxy-3methyl glutaric acid on plasma and low-density lipoprotein cholesterol levels in familial hypercholesterolemia. *Journal of Clinical Pharmacology*, 19:120-126.
- MacRae, W.D. and Towers, G.H.N. (1984). Biological activities of lignans. *Phytochemistry*, 23:1207-1220.
- Matsuda, S., Norimoto, F., Matsumoto, Y., Onba, R., Teramoto, Y., Ohta, N. and Ueda, S. (1994). Solubilization of a novel isoflavone glycoside-hydrolyzing β -glucosidase from *Lactobacillus casei* subsp. rhamnosus IFO 3425. *Journal of Fermentation and Bioengineering*, Osaka, v.77, n.4, p.349-441.
- Mazur, W.M., Fotsis, T., Wähälä, K., Ojala, S., Salakka, A. and Adlercreutz, H. (1996). Isotope-dilution gas-chromatographic mass-spectrometric method for the determination of isoflavonoids, coumestrol and lignans in food samples. *Analytical Biochemistry*, 233:169-180.
- Mazur, W.M. and Adlercreutz, H. (1998). Natural and anthropogenic environmental oestrogens: the scientific basis for risk assessment: naturally occurring oestrogens in food. *Pure and Applied Chemistry*, 70:1759-1776.

- Mazza, G. and Biliaderis, C.G. (1989). Functional properties of flaxseed mucilage. *Journal of Food Science*, 54:1302-1305.
- Mazza, G. and Oomah, B.D. (1995). Flaxseed, dietary fibre, and cyanogens. In: S.C. Cunnane and L.U. Thompson (Eds.), *Flaxseed in Human Nutrition*. AOCS Press, Champaign, IL, pp. 56-81.
- Meagher, L.P., Beecher, G.R., Flanagan, V.P. and Li, B.W. (1999). Isolation and characterization of the lignans, isolariciresinol and pinoresinol, in flaxseed meal. *Journal of Agricultural and Food Chemistry*, 47:173-3180.
- Meagher, L.P. and Beecher, G.R. (2000). Assessment of data on the lignan content of foods. *J Food Comp Anal*, 13:935-947.
- Milder, I.E.J., Arts, I.C.W., Venema, D.P., Lasaroms, J.J.P., Wähälä, K. and Hollman, P.C.H. (2004). Optimization of a liquid chromatography-tandem mass spectrometry method for quantification of the plant lignans secoisolariciresinol, matairesinol, lariciresinol and pinoresinol in foods. *Journal of Agricultural and Food Chemistry*, 52:4643-4651.
- Morton, M.S., Wilcox, G., Wahlqvist, M.L. and Griffiths, K. (1994). Determination of lignans and isoflavonoids in human female plasma following dietary supplementation. *Journal of Endocrinology*, 142:251-59.
- Nesbitt, P.D., Lam, Y. and Thompson, L.U. (1999). Human metabolism of mammalian lignan precursors in raw and processed flaxseed. *American Journal of Clinical Nutrition*, 69:549-555.
- van Niekerk, J.L, Hendriks, T., Gevers Leuven, J.A., Havekes, L. and de Boer, H.H. (1984). The lipid lowering effects of 3-hydroxy-3-methylglutaric acid and bile acid drainage in WHHL rabbits. *Clinical Science*, 67:439-444.
- Nurmi, T., Voutilainen, S., Nyyssönen, K., Adlercreutz, H. and Salonen, J.T. (2003). Liquid chromatography method for plant and mammalian lignans in human urine. *Journal of Chromatography B*, 789:101-110.
- Obermeyer, W.R., Musser, S.M., Betz, J.M., Casey, R.E., Pohland, A.E. and Page, S.W. (1995). Chemical studies of phytoestrogens and related compounds in dietary supplements: Flax and Chaparral. *Proceedings of the Society for Experimental Biology and Medicine*, 208:6-12.
- Oomah, B.D. and Mazza, G. (1993). Flaxseed proteins – A review. *Food Chemistry*, 48:109-114.
- Oomah, B.D., Kenaschuck, E.O. and Mazza, G. (1995). Phenolic acids in

- flaxseed. *Journal of Agricultural and Food Chemistry*, 43:2016-2019.
- Oomah, B.D., Mazza, G. and Kenaschuck, E.O. (1996). Flavonoid content of flaxseed. Influence of cultivar and environment. *Euphytica*, 90:163-167.
- Oomah, B.D. and Mazza, G. (2000). Bioactive components of flaxseed: occurrence and health benefits. In: F. Shahidi and C.T. Ho (Eds), *Phytochemicals and phytopharmaceuticals*. AOCS Press, Champaign, pp. 105-120.
- Orcheson, L.J., Rickard, S.E., Seidl, M.M. and Thompson, L.U. (1998). Flaxseed and its mammalian lignan precursor cause a lengthening or cessation of estrous cycling in rats. *Cancer Letters*, 125:69-76.
- Park, M.S. and Liepa, G.U. (1982). Effects of dietary protein and amino acids on the metabolism of cholesterol-carrying lipoproteins in rats. *Journal of Nutrition*, 112:1892-1898.
- Phipps, W.R., Martini, M.C., Lampe, J.W., Slavin, J.L. and Kurzer, M.S. (1993). Effect of flax seed ingestion on the menstrual cycle. *Journal of Clinical Endocrinology & Metabolism*, 77:1215-1219.
- Pontoh, J. and Low, N.H. (2002). Purification and characterization of beta-glucosidase from honey bees (*Apis mellifera*). *Insect Biochem. Molec. Biol.*, 32:679-690.
- Prasad, K., Gupta, J.B., Kalra, J., Lee, P., Mantha, S.V. and Bharadwaj, B. (1996). Oxidative stress as a mechanism of cardiac failure in chronic volume overload in canine model. *Journal of Molecular and Cellular Cardiology*, 28:375-385.
- Prasad, K. (1997a). Dietary flax seed in prevention of hypercholesterolemic atherosclerosis. *Atherosclerosis*, 132:69-76.
- Prasad, K. (1997b). Hydroxyl radical-scavenging property of secoisolariciresinol diglucoside (SDG) isolated from flax-seed. *Molecular and Cellular Biochemistry*, 168:117-123.
- Prasad, K. (1999). Reduction of serum cholesterol and hypercholesterolemic atherosclerosis by secoisolariciresinol diglucoside isolated from flaxseed. *Circulation*, 99:1355-1362.
- Prasad, K. (2000a). Oxidative stress as a mechanism of diabetes in diabetic BB prone rats: effect of secoisolariciresinol diglucoside (SDG). *Molecular and Cellular Biochemistry*, 209:89-96.
- Prasad, K. (2000b). Antioxidant activity of secoisolariciresinol diglucoside-derived metabolites, secoisolariciresinol, enterodiol, and enterolactone. *International Journal of Angiology*, 9:220-225.

- Prasad, K. (2001). Secoisolariciresinol diglucoside from flaxseed delays the development of type 2 diabetes in Zucker rat. *Journal of Laboratory and Clinical Medicine*, 138:32-39.
- Prasad, K. (2005). Hypocholesterolemic and antiatherosclerotic effect of flax lignan complex isolated from flaxseed. *Atherosclerosis*, 179:269-275.
- Pszczola, D.E. (2000). Genes and diet: the specialized role ingredients may play. *Food Technology*, 54:82-84, 96.
- Rickard, S.E., Orcheson, L.J., Seidl, M.M., Luyengi, L., Fong, H.S. and Thompson, L.U. (1996). Dose-dependent production of mammalian lignans in rats and in vitro from the purified precursor secoisolariciresinol diglucoside in flaxseed. *Journal of Nutrition*, 126:2012-2019.
- Rickard, S.E. and Thompson, L.U. (1997). Health effects of flaxseed mucilage, lignans. *Inform*, 8:860-865.
- Rickard, S.E., Yuan, Y.V. and Thompson, L.U. (2000). Plasma insulin-like growth factor I levels in rats are reduced by dietary supplementation of flaxseed or its lignan secoisolariciresinol diglucoside. *Cancer Letters*, 161:47-55.
- Saarinen, N.M., Smeds, A., Mäkelä, S.I., Ämmälä, J., Hakala, K., Pihlava, J.M., Ryhänen, E.L., Sjöholm, R. and Santti, R. (2002). Structural determinants of plant lignans for the formation of enterolactone in vivo. *Journal of Chromatography B*, 777:311-319.
- Sano, T., Oda, E., Yamashita, T., Shiramasa, H., Ijiri, Y., Yamashita, T. and Yamamoto, J. (2003). Antithrombotic and anti-atherogenic effects of partially defatted flaxseed meal using a laser-induced thrombosis test in apolipoprotein E and low-density lipoprotein receptor deficient mice. *Blood Coagulation and Fibrinolysis*, 14:707-712.
- Schwass, D.E. and Finnley, J.W. (1984). Heat and alkaline damage to proteins: Racemization and lysinoalanine formation. *Journal of Agricultural and Food Chemistry*, 32:1377-1382.
- Schwartz, H. and Sontag, G. (2006). Determination of secoisolariciresinol, lariciresinol and isolariciresinol in plant foods by high performance liquid chromatography coupled with coulometric electrode array detection. *Journal of Chromatography B*, 838:78-85.
- Setchell, K.D.R., Lawson, A.M., Axelson, M. and Adlercreutz, H. (1979). The excretion of two new phenolic compounds during the human menstrual cycle and in pregnancy, endocrinological cancer, ovarian

- function and disease. In: H. Adlercreutz, R.D. Bulbroock, H.J. van der Molen, A. Vermeulen, and S. Sciarra (Eds.), Proceedings of the IX meeting of the international study group for steroid hormones, Rome, December 5-7, 515:207-215.
- Setchell, K.D.R., Lawson, A.M., Borriello, S.P., Harkness, R., Gordon, H., Morgan, D.M.L., Kirk, D.N., Adlercreutz, H. and Anderson, L.C. (1981). Lignan formation in man-microbial involvement and possible roles in relation to cancer. *The Lancet*, 318:4-7.
- Setchell, K.D.R., Lawson, A.M., Borriello, S.P., Adlercreutz, H. and Axelson, M. (1982). Formation of lignans by intestinal microflora. In: R.A. Malt and R.C.N. Williamson (Eds.), *Falk Symposium 31, Colonic Carcinogenesis*. MTP Press Ltd., Lancaster, pp. 93-97.
- Sharma, R., Hudak, M.L., Perszyk, A.A., Premachandra, B.R., Li, H. and Monteiro, C. (2000). Perinatal lethal form of Gaucher's disease presenting with hemosiderosis. *Am J Perinatol*, 17:201-206.
- Shukla, R., Hilaly, A.K. and Moore, K.M. (2003). Process for obtaining lignan. United States patent 20030055227.
- Sicilia, T., Niemeyer, H.B., Honig, D.M. and Metzler, M. (2003). Identification and stereochemical characterization of lignans in flaxseed and pumpkin seeds. *Journal of Agricultural and Food Chemistry*, 51:1181-1188.
- Simopoulos, A.P. (1999). Essential fatty acids in health and chronic disease. *American Journal of Clinical Nutrition*, 70 (suppl): 560S-569S.
- Smeds, A.I., Saarinen, N.M., Hurmerinta, T.T., Penttinen, P.E., Sjöholm, R.E. and Mäkelä, S.I. (2004). Urinary excretion of lignans after administration of isolated plant lignans to rats: the effect of single dose and ten-day exposures. *Journal of Chromatography B*, 813:303-312.
- Smeds, A.I., Hakala, K., Hurmerinta, T.T., Kortela, L., Saarinen, N.M. and Mäkelä, S.I. (2006). Determination of plant and enterolignans in human serum by high-performance liquid chromatography with tandem mass spectrometric detection. *Journal of Pharmaceutical and Biomedical Analysis*, 41:898-905.
- Stitch, S.R., Toumba, J.K., Groen, M.B., Funke, C.W., Leemhuis, J., Vink, J. and Woods, G.F. (1980). Excretion, isolation and structure of a new phenolic constituent of female urine. *Nature*, 287:738-740.
- Sugano, M and Beynen, A.C. (Eds). (1990). *Dietary proteins, cholesterol*

- metabolism and atherosclerosis. New York: Basel.
- Sung, M.K., Lautens, M. and Thompson, L.U. (1998). Mammalian lignans inhibit the growth of estrogen-independent human colon tumor cells. *Anticancer Research*, 18:1405-1408.
- Thompson, L.U., Robb, P., Serraino, M. and Cheung, F. (1991). Mammalian lignan production from various foods. *Nutrition and Cancer*, 16:43-52.
- Thompson, L.U., Rickard, S.E., Cheung, F., Kenaschuk, E.O. and Obermeyer, W.R. (1997). Variability in anticancer lignan levels in Flaxseed. *Nutrition and Cancer*, 27:26-30.
- Thompson, L.U. (1998). Experimental studies on lignans and Cancer. *Bailliere's Clinical Endocrinology and Metabolism*, 12:691-705.
- Wang, C., Mäkelä, S.I., Hase, T., Adlercreutz, H. and Kurzer, M.S. (1994). Lignans and flavonoids inhibit aromatase enzyme in human preadipocyte. *Journal of Steroid Biochemistry and Molecular Biology*, 50:205-212.
- Wang, L.Q. (2002). Mammalian phytoestrogens: enterodiol and enterolactone. *Journal of Chromatography*, 777:289-309.
- Westcott, N.D. and Muir, A.D. (1996). Process for extracting and purifying lignans and cinnamic acid derivatives from flaxseed. PCT Patent WO9630468A2.
- Xu, X., Harris, K.S., Wang, H.J., Murphy, P.A. and Hendrich, S. (1995). Bioavailability of soybean isoflavones depends upon gut microflora in women. *Journal of Nutrition*, 125:2307-2315.
- Yamauchi, F. and Suetsuna, K. (1993). Immunological effects of dietary peptide derived from soybean protein. *J. Nutr. Biochem.* 4:450-457.
- Yu, H.L., Xu, J.H., Lu, W.Y. and Lin, G.Q. (2007). Identification, purification and characterization of β -glucosidase from apple seed as a novel catalyst for synthesis of O-glucosides. *Enzyme Microb Technol*, 40:354-361.
- Yuan, J.P., Li, X., Xu, S.P., Wang, J.H. and Liu, X. (2008). Hydrolysis kinetics of secoisolariciresinol diglucoside oligomers from flaxseed. *J Agric Food Chem*, 56(21):10041-7.