

捌、參考文獻

王霞、高麗娟及林炳昌（2005）。表沒食子兒茶素沒食子酸酯(EGCG)的分離與製備。食品科學，26：242-246。

仇燕嶸、李楠、韓國柱及郭永學（2006）。表沒食子兒茶素沒食子酸酯的研究進展。中草藥，37：303-306。

安田英之（1992）。兒茶素類的消臭作用及利用。食品工業，35：28-33。

何小解、盧向羊、易著文、方俊、田云、黨西強及莫雙紅（2005）。模擬人體胃腸道環境對兒茶素穩定性的影響。湖南農業大學學報（自然科學版），31：527-529。

李昂（2001）。奈米級顆粒在藥物輸遞的應用。化工資訊，10：44-55。

沈生榮（1993）。茶多酚的穩定性。茶葉，2：27-31。

林佳姣及林睿哲（2002）。幾丁質與幾丁聚糖於生醫材料之應用與特性。化工技術，10：228-237。

沈家弘（2004）。不同去乙醯度幾丁聚糖乳化性質之比較及應用於沙拉醬之研究。私立東海大學食品科學研究所碩士論文。

河內二郎（1987）。茶葉抽出物消臭的發展。食品工業，22：34-35。

柯建邦（2006）。不同去乙醯度膠態幾丁聚糖乳化性質之比較及應用於沙拉醬之研究。私立東海大學食品科學系研究所碩士論文。

徐世昌（2001）。生物性高分子幾丁質與幾丁聚糖之介紹與應用。化工資訊，2：36-45。

徐志泉、易著文及何小解（2006）。表食子酸酯對大鼠腎衰的治療研究進展。現代醫藥衛生，22：471-472。

孫勇、李華佳、辛志宏及胡秋輝（2006）。奈米食品的活性與安全性研究。食品科學，27：936-939。

屠幼英 (1994)。茶提取物理化學特性的研究。食品科學，3；11-14。

莊仲揚 (2002)。幾丁聚醣於生醫產業上的應用。化工資訊，4：46-50。

莊景光 (2004)。離子鍵結型奈米微粒製備與其小腸上皮細胞滲透能力之探討。國立清華大學化學工程研究所碩士論文。

郭麗雪 (2003)。幾丁聚醣-果膠-海藻酸鈉複合膠包覆兒茶素其釋放特性之研究。國立台灣海洋大學食品科學研究所碩士論文。

葉安義 (2004)。奈米科技與食品。科技發展，394：44-49。

賴淑琪 (1979)。水產廢棄物蝦、蟹外殼之高度利用。食品工業，11：23-28。

陳英玲 (2005)。茶葉的保健功效。科學發展，391：66-73。

陳美惠、莊淑惠及吳志律 (2000)。幾丁聚醣的物化特性。食品工業，31：1-6。

謝馥如 (2007)。奈米食品安全性與生體利用率。食品工業，39：35-45。

顏子揚 (2007)。茶飲料中兒茶素對微生物的抑菌效應。食品工業，39：39-47。

鄭育奇 (2008)。生物高分子材料在機能性食品之製造應用。食品工業，40：45-53。

Agnihotri S. A., Mallikarjuna N. N. and Aminabhavi T. M. (2004). Recent advances on chitosan-based micro- and nanoparticles in drug delivery. *J. Control. Release.*, 10 : 5-28.

Agnihotri S. A. and Aminabhavi T. M. (2004). Contralled release of clozapine through chitosan micropaticles prepared by a novel method. *J. Control. Release.*, 96:245-259.
Ahmad N., Gupta S. and Mukhtar H. (2000). Green tea polyphenol pigallocatechin allate differentially modulates nuclear factor κB in cancer ells versusnormal cells. *Arch. Biochem. Biophys.*, 376 : 338-346.

Akbuga J. and Durmaz G. (1994). Preparation and evaluation of crosslinked chitosan microspheres containing furosemide. *Int. J. Pharm.*, 11: 217-222.

Allan C. R. and Hadiger L. A. (1979). The fungicidal effect of chitosan on fungi of varying cell wall composition. *Exp. Mycol.*, 3 : 285-287.

Allan G. G. and Peyron M. (1995). Molecular weight manipulation of chitosan I: kinetic of depolymerization by nitrous acid. *Carbohydr. Res.*, 27 : 257-272 .

Ballard S. T., Hunter J. H. and Taylor A.(1995). Regulation of tight-junction permeability during nutrient absorption across the intestinal epithelium. *Annu. Rev. Nutr.*, 15 : 35-55.

Bandyopadhyay D., Chatterjee T. K., Dasgupta A., Lourduraja J. and Dastidar S. G. (2005). In vitro and in vivo antimicrobial action of tea: the commonest beverage of Asia. *Biol. Pharm. Bull.*, 28 : 2125-2127.

Berthod A. and Kreuter J. (1996). Chitosan microspheres-improved acid stability and change in physicochemical properties by cross-linking. *Proc. Int. Symp. Control. Release Bioact. Mater.*, 23 : 369-370.

Besse J., Besse L. and Alphonsr M. (2006). Impregnated powder improving bioavailability and/or the solubility and method of production. US Patent 2006018269 A1.

Bhumkar D. R. and Pokharkar V. B. (2006). Studies on effect of pH on cross-linking of chitosan with sodium tripolyphosphate: a technical note. *AAPS PharmSciTech*. 7 : E50.

Cabrera C., Artacho R. and Giménez R. (2006). Beneficial Effects of Green Tea—A Review. *J. Am. Coll. Nutr.*, 25 : 79-99.

Calvo P., Remunan-Lopez C., Vila-Jato J. L. and Alonso M. J. (1997). Novel hydrophilic chitosan-polyethylene oxide nanoparticles as protein carriers. *J. Appl. Polym. Sci.*, 63 : 125-132.

Catterall F., King L. J., Clifford M. N. and Ioannides C. (2003). Bioavailability of dietary doses of ^3H -labelled tea antioxidants (+)-catechin and (-)-epicatechin in rat. *Xenobiotica*, 33 : 743–753.

Chan M. M., Ho C. T. and Huang H. I. (1995). Effects of three dietary phytochemicals

from tea, rosemary and turmeric on inflammation-induced nitrite production. *Cancer Lett.*, 96 : 23-29.

Chandy T. and Sharma C. P. (1990). Chitosan-as a biomaterial. *Biomat. Art. Cells Art. Org.*, 18 : 1–24.

Cheng J. Y., Mao Q. L., Kuang F. C., Lu Y. and Shi Z. P. (2007). Research advances in health and pharmacological functions of tea catechins. *Tea Commun.*, 34 : 14-18.

Collares-Buzato C. B., Jepson M. A. Simmons N. L. and Hirst B. H. (1998).Increased tyrosine phosphorylation causes redistribution of adherens junction and tight junction proteins and perturbs paracellular barrier function in MDCK epithelia. *Eur. J. Cell Biol.*, 76 : 85–92.

Deng Q. Y., Zhou C. R. and Luo B. H. (2006). Preparation and characterization of chitosan nanoparticles containing lysozyme. *Pharm. Biol.*, 44 : 336-342.

Desai M. P., Labhsetwar V., Walter E., Levy R. J. and Amidon G. L. (1997). The mechanism of uptake of biodegradable microparticles in Caco-2 cells is size dependent. *Pharm. Res.*, 14: 1568-73.

Fang J. Y., Lee W. R., Shen S. C. and Huang Y. L. (2006). Effect of liposome encapsulation of tea catechins on their accumulation in basal cell carcinomas. *J. Dermatol. Sci.*, 42 : 101-109.

Feng S. S. and Chien S. (2003). Chemotherapeutic engineering: Application and further development of chemical engineering principles for chemotherapy of cancer and other diseases. *Chem. Eng. Sci.*, 58 : 4087-4114.

Fernandez-Urrusuno R., Calvo P., Remunan-Lopez J. L. and Alonso M. J. (1999). Enhancement of nasal absorption of insulin using chitosan nanoparticles. *Pharm. Res.*, 16 : 1576-1581.

Furuse M., Itoh M., Hirase T., Nagafuchi A., Yonemura S. and Tsukita S. (1994). Direct association of occluding with ZO-1 and its possible involvement in the localization of occluding at tight junctions. *J. Cell Biol.*, 127 : 1617–26.

Gai Y., Anavy N. D. and Chow H. S. S. (2002).Contribution of presystemic hepatic

extraction to the low oral bioavailability of green catechins in rat. *Drug Metab. Dispos.*, 30 : 1246-1249.

Gan Q., Wang T., Cochrane C. and McCarron P. (2005). Modulation of surface change, particle size and morphological properties of chitosan-TPP nanoparticles intended for gene delivery. *Colloids Surf. B Biointerfaces*, 44 : 65-73.

George M. and Abraham T. E. (2006). Polyionic hydrocolloid for the intestinal delivery of protein drug: Alginate and chitosan- a review. *J. Control. release*, 114 : 1-14.

Guan R. F., Qian B., Ye X. Q. and Hao Y. B. (2006). Recent study of nanotechnology in food science. *Food Sci.*, 27 : 270-273.

He P., Davis S. S. and Illum L. (1998). In vitro evaluation of the mucoadhesive properties of chitosan microspheres. *Int. J. Pharm.*, 166 : 75-88.

He P., Davis, S. S. and Illum, L. (1999). Chitosan microspheres prepared by spray drying. *Int. J. Pharm.*, 187 : 53-65.

Hoekstra A., Stuszczyk H. and Kivekas O. (1998). Percutaneous microcrystalline chitosan application for sealing arterial puncture sites. *Biomaterials*, 19 : 1467-1471.

Hollman P. C. H. (2004). Absorption, bioavailability and metabolism of flavonoids. *Pharma. Biol.*, 42: 74-83.

Hsieh F. M., Huang C., Lin T. F., Chen Y. M. and Lin J. C. (2008). Study of sodium tripolyphosphate-crosslinked chitosan beads entrapped with *Pseudomonas putida* for phenol degradation. *Process Biochemistry*, 43 : 83-92.

Hu B., Pan C., Sun Y., Hou Z., Ye H., Hu B. and Zeng X. (2008). Optimization of fabrication parameters to produce chitosan-tripolyphosphate nanoparticles for delivery of tea catechins. *J. Agric. Food Chem.*, 56 : 7451-7458.

Huang M., Khor E. and Lim L. Y. (2004). Uptake and cytotoxicity of chitosan molecules and nanoparticles: effect of molecular weight and degree of deacetylation. *Pharm. Res.*, 21 : 344-353.

Ishigami T. (1991). Proceedings of the international symposium on Tea Science. Shizuoka, Japan. 248-252.

Janes K. A., Fresneau M. P., Marazuela A., Fabra A. and Alonso M. J. (2001). Chitosan nanoparticles and delivery systems for doxorubicin. *J. Control. Rel.*, 73 : 255-267.

Janes K. A. and Alonso M. J. (2003). Depolymerized chitosan nanoparticles for protein delivery: preparation and characterization. *J. Appl. Polym. Sci.*, 88 : 2769-2776.

Jeon Y. J., Shahidi F. and Kim S. K. (2000). Preparation of chitin and chitosan oligomers and their application in physiological functional foods. *Food Rev. Int.*, 16 : 159-176.

Kawashima Y., Handa T., Takenaka H., Lin S. Y. and Ando Y. (1985). Novel method for the preparation of controlled-release theophylline granules coated with a polyelectrolyte complex of sodium polyphosphate-chitosan. *J. Pharm. Sci.*, 74 : 264-268.

Knaul J. Z., Hudosn S. M. and Creber K. A. M. (1999). Improved mechanical properties of chitosan fibers. *J. Appl. Polym. Sci.*, 72 : 1721-1731.

Ko J. A., Park H. J., Hwang S. J., Park J. B. and Lee J. S. (2002). Preparation and characterization of chitosan microparticles intended for controlled drug delivery. *Int. J. Pharm.*, 249 : 165-174.

Koo M. W. L. and Cho C. H. (2004). Pharmacological effect of green tea on the gastrointestinal system. *Eur. J. Pharm.*, 500: 177-185.

Kujala T. S., Loponen J. M., Klika K. D. and Pihlaja K. (2000). Phenolics and betacyanins in red beetroot (*Beta vulgaris*) root: distribution and effect of cold storage on the content of total phe-nolics and three individual compounds. *J. Agric. Food Chem.*, 48 : 5338-5342.

Leong Y. S. and Candau F. (1982). Inverse microemulsion polymerization. *J. Phys. Chem.*, 86 : 2269-2271.

Lin Y. H., Chung C. K., Chen C. T., Liang H. F., Chen S. C. and Sung H. W. (2005). Preparation of nanoparticles composed of chitosan/poly- γ -glutamic acid and evaluation of their permeability through caco-2 cells. *Biomacromolecules*, 6 : 1104–1112.

Lin Y. H., Chen C. T., Liang H. F., Kulkarni A. R., Lee P. W., Chen C. H. and Sung W. H. (2007). Novel nanoparticles for oral insulin delivery via the paracellular pathway. *Nanotechnology*, 18 : 105102.

Lin Y., Tang J., Chen X. and Xin H. (2005). A templating route to nanoporous chitosan materials. *Carbohydr. Res.*, 340 : 2816-2820.

Liu H., Huang K. and Xu H. (2006). Recent research on biological safety of nanometer-scale materials. *Chem. Indust. Engin. Prog.*, 25 : 1040-1044.

Mi F. L., Shyu S. S., Lee S. T. and Wong T. B. (1999a). Kinetic study of chitosan-tripolyphosphate complex reaction and acid-resistive properties of the chitosan-tripolyphosphate gel beads prepared by in-liquid curing method. *J. Polym. Sci. Part B. Polymer physics.*, 37 : 1551-1564.

Mi F. L., Shyu S. S., Kuan C. Y., Lee S. T., Lu K. T. and Jang S. F. (1999 b). Chitosan-tripolyphosphate complexation for the preparation of gel beads and controlled release of anticancer drug. I. effect of phosphorous polyelectrolyte complex and enzymatic hydrolysis of polymer. *J. Appl. Polym. Sci.*, 74 : 1868-1879.

Mohanraj V. J. and Chen Y. (2006). Nanoparticle- A Review. *Trop. J. Pharm. Res.*, 5 : 561-573.

Mullin M. J., Kampherstein J. A., Laughlin K. V., Cheryl E. K., Clarkin R., Miller D., Szallasi Z., Kachar B., Soler A. P. and Rosson D. (1998). Overexpression of protein kinase C- δ increases tight junction permeability in LLC-PK1 epithelia. *Am. J. Physiol.*, 275 : 544–554.

Muzzarelli R. A. A. and Rocchetti R. (1985). Determination of the degree of acetylation of chitosans by first derivative ultraviolet spectrophotometry. *Carbohydr. Polymers*, 5 : 461-472.

Nishimura, K., Nishimura, S., Seo, H., Nishi, N., Tokura, S. and Azuma, I. (1986).

Macrophage activation with multiporous beads prepared from partially deacetylated chitin. *J. Biomed. Mater. Res.*, 20 : 1359-1372

Nishimura K., Nishimura S., Seo H., Nishi N., Tokura S. and Azuma I. (1987). Effect of multiporous microspheres derived from chitin on the activation of mouse peritoneal macrophages. *Vaccine*, 5 : 136-140.

Nishimura K., Nishimura S., Nishi N., Saiki I., Tokura S. and Azuma I. (1992).

Advances in Chitin and Chitosan. pp. 477-483, Brine, C. J., Sandford, P. A. and Zikakis, J. P. (ed.), Elsevier Appl. Sci., N. Y.

Okamoto Y., Shibasaki K., Minami S., Matsuhashi A., Tanioka S. and Shigemasa Y. (1995). Evaluation of chitin and chitosan on open wound healing in dogs. *J. Vet. Med. Sci.*, 57 : 851-854.

Pan Y., Li Y. J., Zhao H.Y., Zheng J. M., Xu H., Wei G., Hao J. S. and Cui F. D. (2002). Bioadhesive polysaccharide in protein delivery system: chitosan nanoparticles improve the intestinal absorption of insulin in vivo. *Int. J. Pharm.*, 249 : 139-147.

Pantaleone D., Yalpani M. and Scollar M. (1992). Unusual susceptibility of chitosan to enzymic hydrolysis. *Carbohydr. Res.* 237 : 325-332.

Panyam J. and Labhsetwar V. (2003). Biodegradable nanoparticles for drug and gene delivery to cells and tissue. *Adv. Drug. Deliv. Rev.*, 55 : 329-47.

Papadimitriou S., Bikaris D., Avgoustakis K., Karavas E. and Georgarakis M. (2008). Chitosan nanoparticles loaded with dorzolamide and pramipexole. *Carbohydr. Polymers*, 73 : 44-54.

Peluso G., Petillo O., Ranieri M., Santin M., Ambrosio L., Calabro D., Avallone B. and Balsamo G. (1994). Chitosan-mediated stimulation of macrophage function. *Biomaterials*, 15 : 1215-1220.

Qi L., Xu Z., Jiang X., Hu C. and Zou X. (2004). Preparation and antibacterial activity of chitosan nanoparticles. *Carbohydr. Res.* 339 : 2693-2700.

Qi L., Xu Z. and Chen M. (2006). In vitro and in vivo suppression of hepatocellular carcinoma growth by chitosan nanoparticles. *Eur. J. Cancer*, 43 : 184-193.

Qin C., Zhou B., Zeng L., Zhang Z., Liu L., Du Y. and Xiao L. (2004). The physicochemical properties and antitumor activity of cellulase-treated chitosan. *Food chem.*, 84 : 107-115.

Ribeiro A. J., Neufeld R. J., Arnaud P. and Chaumeil J. C. (1999). Microencapsulation of lipophilic drug in chitosan-coated alginate microsphere. *Int. J. Pharm.* 187 : 115-123.

Rinaudo M.(2006). Chitin and chitosan: Properties and applications. *Prog. Polym. Sci.*, 31 : 603-632.

Ruiz-Herrera J.(1978). The distribution quantitative importance of chitin in fungi. In: chitin and chitosan, Proceedings of the First International Conference on Chitin and Chitosan. Muzzarelli, R. A. A. and Pariser, E. R. Eds.MIT Sea Grant Program, Cambridge, Mass. P.11.

Soane R. J., Frier M., Perkins A. C., Jones N. S., Davis S. S. and Illum L. (1999). Evaluation of the clearance characteristics of bioadhesive systems in humans. *Int. J. Pharm.*, 178 : 55-56.

Soppimath K. S., Aminabhavi T. M., Kulkarni A. R. and Rudzinski W. E. (2001). Biodegradable polymeric nanoparticles as drug delivery devices. *J. Control. Release*, 70 : 1-20.

Struszczyk H., Wawro D. and Niekraszewicz A. (1991). Biodegradability of chitosan fibres. In: C.J. Brine, P.A. Sandford and J.P. Zikakis, Editors, *Advances in Chitin and Chitosan*, Elsevier Applied Science, London , 580–585.

Suganuma M ., Okabe S. and Sueoka E. (1996). A new process of cancer prevention mediated through inhibition of tumor necrosis factor α expression. *Cancer Res.*, 56 : 3711-3715.

Tai Y. H., Flick J., Levine S. A., Madara J. L., Sharp G. W. and Donowitz M. (1996). Regulation of tight junction resistance in T84 monolayers by elevation in intracellular Ca^{2+} a protein kinase C effect. *J. Membr. Biol.*, 149 : 71–79.

Thanou M., Verhoef J. C. and Junginger H. E. (2001). Oral drug absorption enhancement by chitosan and its derivatives. *Adv. drug deliv. rev.*, 52 : 117-126.

Toei K. and Kohara T. (1976). A conductometric method for colloid titrations. *Analytica. Chimica. Acta.*, 83 : 59-65.

Tokumitsu H., Ichikawa H. and Fukumori Y. (1999). Chitosan-gadopentetic acid complex nanoparticles for gadolinium neutron capture therapy of cancer : preparation by novel emulsion-droplet coalescence technique and characterization. *Pharm. Res.*, 16 : 1830-1835.

Varilek G. W., Yang F., Lee E. Y., deVilliers W. J. S., Zhong J., Oz H. S., Westberry K. F. and McClain C. J. (2001). Green tea polyphenol extract attenuates inflammation in interleukin-2-deficient mice, a model of autoimmunity. *J. Nutr.*, 131 : 2034-2039.

Vila A., Sanchez A., Janes K., Behrens I., Kissel T., Vila-Jato J. L. and Alonso M. J. (2004). Low molecular weight chitosan nanoparticles as new carriers for nasal vaccine delivery in mice. *Eur. J. Pharm. Biopharm.*, 57 : 123-131.

Ward P. D., Tippin T. K. and Thakker D. R. (2000). Enhancing paracellular permeability by modulating epithelial tight junctions. *Pharm. Sci. Technolo. Today*, 3 : 346-358.

Wang W., BO S., Li S. and Qin W. (1991). Determination of the Mark-Houwink equation for chitosans with different degrees of deacetylation. *Int. J. Biol. Macromol.*, 13 : 281-285.

Wu G. J. and Tsai G. J. (2004). Cellulase degradation of shrimp chitosan for the preparation of a water-soluble hydrolysate with immunoactivity. *Fish. Sci.*, 70 : 1113–1120.

Wu Y., Yang W., Wang C., Hu J. and Fu S. (2005). Chitosan nanoparticles as a novel delivery system for ammonium glycyrrhizinate. *Int. J. Pharm.*, 295 : 235-245.

Xu Y. and Du Y. (2003). Effect of molecular structure of chitosan: protein delivery properties of chitosan nanoparticles. *Int. J. Pharm.*, 250 : 215-226.

Yalpani M. and Pantaleone D. (1994). An examination of the unusual susceptibilities of aminoglycans to enzymatic hydrolysis. *Carbohydr. Res.*, 256 : 159-175.

Yamamoto T., Harada N., Kano K., Taya S., Canaani E., Matsuura Y., Mizoguchi A., Ide C. and Kaibuchi K. (1997). The ras target AF-6 interacts with ZO-1 and serves as a peripheral component of tight junctions in epithelial cells. *J. Cell. Biol.*, 139 : 785–95.

Yanagawa Y., Yamamoto Y., Hara Y. and Shimamura T. (2003). A Combination effect of epigallocatechin gallate, a major compound of green tea catechins, with antibiotics on helicobacter pylori growth in Vitro. *Curr. Microbiol.*, 47 : 244-249.

Yang T. T. C. and Koo M. W. L. (1997). Hypocholesterolemic effects of Chinese tea .*Pharma. Res.*, 35: 505-512.

Yu J. H., Du X. M. and Li X. H. (1999). Bland films of chitosan/gelation. *J. Wuhan Univ.*, 45 : 440-444.

Zeng L., Qin C., Wang W., Chi W. and Li W. (2007). Aborption and distribution of chitosan in mice after oral administration. *Carbohydr. polymers*, 71 : 435-440.

Zhang M., Li X. H., Gong Y. D., Zhao N. M. and Zhang X. F. (2002) . Properties and biocompatibility of chitosan films modified by blending with PEG. *Biomaterials*, 23 : 2641-2648.

Zhang L. and Kosaraju S. L. (2007). Biopolymeric delivery system for controlled release of polyphenolic antioxidants. *Eur. Polym. J.*, 43 : 2956-2966.