

參考資料

- [1] M. L. Grayeski, Coumarin derivatizing agents for carboxylic acid detection using peroxyoxalate chemiluminescence with liquid chromatography, *Anal. Chem.*, **1987**, 59, 1203-1206.
- [2] A. Townshend, Solution chemiluminescence—some recent analytical developments, *Analyst*, **1990**, 115, 495-500.
- [3] W. R. G. Baeyens, B. L. Ling, K. Imai, A. C. Calokerinos, S. G. Schulman, Chemiluminescence detection in capillary electrophoresis, *J. Microcol. Sep.*, **1994**, 6, 195-206.
- [4] J. W. Birks, Chemiluminescence and photochemical reactions detection in chromatography, VCH publishers Inc., New York, **1989**.
- [5] L. J. Kricka, G. H. G. Thorpe, Chemiluminescent and bioluminescent methods in analytical chemistry. A review, *Analyst*, **1983**, 108, 1274-1296.
- [6] K. Robard, P. J. Worsfold, Analytical applications of liquid-phase chemiluminescence, *Anal. Chim. Acta*, **1992**, 266, 147-173.
- [7] P. J. M. Kwakman, U. A. T. Brinkman, Peroxyoxalate chemiluminescence detection in liquid chromatography., *Anal. Chim. Acta*, **1992**, 266, 175-192.
- [8] A. W. Knight, A review of recent trends in analytical applications of electrogenerated chemiluminescence, *Trends Anal. Chem.*, **1999**, 18, 47-62.
- [9] R. T. Dufford, D. Nightingale, L. W. Gaddnm, Luminescence of grignard compounds in electricand magnetic fields, and related electrical phenomena, *J. Am. Chem. Soc.*, **1927** , 49, 1858-1864.
- [10] N. Harvey, Luminescence during electrolysis, *J. Phys. Chem.*, **1929**, 33, 1456-1459.
- [11] M. M. Richter, Electrochemiluminescence (ECL), *Chem. Rev.*, **2004**,

104, 3003-3036.

- [12] F. H. Burstall, Optical activity dependent on co-ordinated bivalent ruthenium, *J. Chem. Soc.*, **1936**, 58, 173-175.
- [13] J. Steigman, N. Birnbaum, S. M. Edmonds, Ruthenium dipyriddy-new oxidimetric indicator, *Ind. Eng. Chem. Anal. Ed.*, **1942**, 14, 30.
- [14] J. P. Paris, W. W. Brandt, Charge transfer luminescence of a ruthenium(II) chelate, *J. Am. Chem. Soc.*, **1959**, 81, 5001-5002.
- [15] H. Veening, W. W. Brandt, Fluorescimetric determination of ruthenium, *Anal. Chem.*, **1960**, 32, 1426-1428.
- [16] D. M. Hercules, F. E. Lytle, Chemiluminescence from reduction reactions, *J. Am. Chem. Soc.*, **1966**, 88, 4745-4746.
- [17] I. Rubinstein, A. J. Bard, Electrogenerated chemiluminescence. 37. Aqueous ecl systems based on tris(2,2'-bipyridine)ruthenium(2+) and oxalate or organic acids, *J. Am. Chem. Soc.*, **1981**, 103, 512-516.
- [18] O. Shinozaki, N. Kunezaki, S. Yamazaki, T. Tanimura, Chemiluminescent method for aliphatic tertiary amines using $\text{Ru}(\text{bipy})_3^{2+}$ - S_2O_8 reagent, *Chromatographia*, **1992**, 13, 275-276.
- [19] S. Yamazaki, R. Chiba, K. Uchikura, T. Tanimura, Simultaneous determination of primary, secondary and tertiary amines by postcolumn reaction, *Chromatographia*, **1995**, 16, 322-323.
- [20] S. Yamazaki, K. Ozaki, K. Saito, T. Tanimura, Enantiomeric separation of β -amino alcohols with a primary or tertiary amine moiety by reversed phase liquid chromatography with a chiral mobile phase containing copper (II) and (*R*)-mandelic acid, *J. High Resolut. Chromatogr.*, **1995**, 18, 68-70.
- [21] I. N. Demas, G. A. Crosby, On the multiplicity of the emitting state of ruthenium(II) complexes, *J. Mol. Spectrosc.*, **1968**, 26, 72-77.
- [22] F. E. Lytle, D. M. Hercules, Luminescence of tris(2,2'-bipyridine)ruthenium(II) dichloride, *J. Am. Chem. Soc.*, **1969**,

- 91, 253-257.
- [23] J. N. Demas, G. A. Crosby, Quantum efficiencies on transition metal complexes. II. Charge-transfer luminescence, *J. Am. Chem. Soc.*, **1971**, *93*, 2841-2847.
- [24] R. D. Gerardi, N. W. Barnett, S. W. Lewis, Analytical applications of tris(2,2'-bipyridyl)ruthenium(III) as a chemiluminescent reagent, *Anal. Chim. Acta*, **1999**, *378*, 1-41.
- [25] A. W. Knight, G. M. Greenway, Relationship between structural attributes and observed electrogenerated chemiluminescence (ECL) activity of tertiary amines as potential analytes for the tris(2,2'-bipyridine)ruthenium(II) ECL reaction. A review, *Analyst*, **1996**, *121*, 101R-106R.
- [26] J. B. Noffsinger, N. D. Danielson, Generation of chemiluminescence upon reaction of aliphatic amines with tris(2,2'-bipyridine)ruthenium(III), *Anal. Chem.*, **1987**, *59*, 865-868.
- [27] J. B. Noffsinger, N. D. Danielson, Liquid chromatography of aliphatic trialkylamines with post-column chemiluminescent detection using tris(2,2'-bipyridine) ruthenium(III), *J. Chromatogr.*, **1987**, *387*, 520-524.
- [28] K. Uchikura, M. Kirisawa, Generation of chemiluminescence upon reaction of alicyclic tertiary amines with tris(2, 2'-bipyridine) ruthenium(II) using flow electrochemical reactor, *Anal. Sci.*, **1991**, *7*, 803-804.
- [29] T. M. Downey, T. A. Nieman, Chemiluminescence detection using regenerable tris(2,2'-bipyridyl)ruthenium(II) immobilized in Nafion, *Anal. Chem.*, **1992**, *64*, 261-268.
- [30] K. Uchikura, M. Kirisawa, A. Sugii, Electrochemiluminescence detection of primary amines using tris(bipyridine)ruthenium(III) after derivatization with divinylsulfone, *Anal. Sci.*, **1993**, *9*, 121-123.
- [31] S. N. Brune, D. R. Bobbitt, Role of electron-donating/withdrawing

- character, pH, and stoichiometry on the chemiluminescent reaction of tris(2,2'-bipyridyl)ruthenium(III) with amino acids, *Anal. Chem.*, **1992**, *64*, 166-170.
- [32] M. Shirao, R. Furuta, S. Suzuki, H. Nakazawa, S. Fujita, T. Maruyama, Chemiluminescence detection of amino acids, peptides, and proteins using tris-2,2'-bipyridine ruthenium (III), *Anal. Lett.*, **1990**, *23*, 195-210.
- [33] S. N. Brune, D. R. Bobbitt, Effect of pH on the reaction of tris tris(2,2'-bipyridyl)ruthenium(III) with amino-acids : Implications for their detection, *Talanta*, **1991**, *38*, 419-424.
- [34] K. Uchikura, M. Kirisawa, Chemiluminescence of Tryptophan with electrogenerated tris(2,2'-bipyridine)ruthenium(III), *Chem. Lett.*, **1991**, *8*, 1373-1376.
- [35] K. Uchikura, M. Kirisawa, Notes electrochemiluminescence determination of d-, l-tryptophan using ligand-exchange high-performance liquid chromatography, *Anal. Sci.*, **1991**, *7*, 971-973.
- [36] W. Y. Lee, T. A. Nieman, Determination of dansyl amino acids using tris(2,2'-bipyridine)ruthenium(III) chemiluminescence for post-column reaction detection in high-performance liquid chromatography, *J. Chromatogr. A*, **1994**, *659*, 111-118.
- [37] D. R. Skotty, T. A. Nieman, Determination of oxalate in urine and plasma using reversed-phase ion-pair high-performance liquid chromatography with tris(2,2'-bipyridyl)ruthenium(II)-electrogenerated chemiluminescence detection, *J. Chromatogr. B*, **1995**, *665*, 27-36.
- [38] N. Egashira, H. Kumasako, K. Ohga, Fabrication of a Fiber-Optic-Based Electrochemiluminescence sensor and its application to the determination of oxalate, *Anal. Sci.*, **1990**, *6*, 903-904.
- [39] G. P. Jirka, T. A. Nieman, Modulated potential electrogenerated chemiluminescence of luminol and $\text{Ru}(\text{bpy})_3^{2+}$, *Mikrochim. Acta*, **1994**, *113*, 339-347

- [40] I. Rubinstein, A. J. Bard, Electrogenerated chemiluminescent determination of oxalate, *Anal. Chem.* **1983**, *55*, 1580-1582.
- [41] J. A. Holeman, N. D. Danielson, Liquid chromatography of antihistamines using post-column tris(2,2'-bipyridine) ruthenium(III) chemiluminescence detection, *J. Chromatogr. A*, **1994**, *679*, 277-284.
- [42] A. F. Martin, T. A. Nieman, Glucose quantitation using an immobilized glucose dehydrogenase enzyme reactor and a tris(2,2'-bipyridyl) ruthenium(II) chemiluminescent sensor, *Anal. Chim. Acta*, **1993**, *281*, 475-481.
- [43] G. A. Forbes, T. A. Nieman, J. V. Sweedler, On-line electrogenerated Ru(bpy)₃³⁺ chemiluminescent detection of β-blockers separated with capillary electrophoresis, *Anal. Chim. Acta*, **1997**, *347*, 289-293.
- [44] D. R. Bobbit, W. A. Jackson, H. D. Hendricloson, Chemiluminescent detection of amines and amino acids using in situ generated Ru(bpy)₃³⁺ following separation by capillary electrophoresis, *Talanta*, **1998**, *46*, 565-572.
- [45] M. T. Chiang, C. W. Whang, Tris(2,2'-bipyridyl)ruthenium(III)-based electrochemiluminescence detector with indium/tin oxide working electrode for capillary electrophoresis, *J. Chromatogr. A*, **2001**, *934*, 59-66.
- [46] K. A. Fahrnich, M. Pravda, G. G^l. Guilbault, Recent applications of electrogenerated chemiluminescence in chemical analysis, *Talanta*, **2001**, *54*, 531-559.
- [47] I. Rubinstein, C.R. Martin, A.J. Bard, Electrogenerated chemiluminescent determination of oxalate, *Anal. Chem.*, **1983**, *55*, 1580-1582.
- [48] J.B. Noffsinger, N.D. Danielson, Generation of chemiluminescence upon reaction of aliphatic amines with tris(2,2'-bipyridine)ruthenium(III), *Anal. Chem.* **1987**, *59*, 865-868.

- [49] K. Uchikura, Determination of oxalate by FIA with electrogenerated chemiluminescence detection, *Bunseki Kagaku*, **1990**, *39*, 323-326.
- [50] N. Egashira, H. Kumasako, K. Ohga, Fabrication of a fiber-optic-based electrochemiluminescence sensor and its application to the determination of oxalate, *Anal. Sci.*, **1990**, *6*, 903-904.
- [51] N. Egashira, H. Kumasako, Y. Kurauchi, K. Ohga, Determination of oxalate in vegetables with a fiber-optic electrochemiluminescence sensor, *Anal. Sci.*, **1992**, *8*, 713-714.
- [52] T. M. Downey, T. A. Nieman, Chemiluminescence detection using regenerable tris(2,2'-bipyridyl)ruthenium(II) immobilized in Nafion, *Anal. Chem.*, **1992**, *64*, 261-268.
- [53] N. Egashira, H. Kumasako, Y. Kurauchi, K. Ohga, Determination of oxalate with fiber-optic electrochemiluminescence sensors., *Proc. Electrochem. Soc.*, **1993**, *7*, 674-679.
- [54] D. J. Walton, S. S. Phull, D. M. Bates, J. P. Lorimer, T. J. Mason, Sonochemical enhancement of electrochemiluminescence. *Ultrasonics*, **1992**, *30*, 186-191.
- [55] G.P. Jirka, T.A. Nieman, Modulated potential electrogenerated chemiluminescence of luminol and $\text{Ru}(\text{bpy})_3^{2+}$. *Mikrochim. Acta* , **1994**, *113*, 339-347.
- [56] N. W. Barnett, T. A. Bowser, R. A. Russell, Determination of oxalate in alumina process liquors by ion chromatography with post column chemiluminescence detection, *Anal. Proc.*, **1995**, *32*, 57-59.
- [57] D. R. Skotty, T. A. Nieman, Determination of oxalate in urine and plasma using reversed-phase ion-pair high-performance liquid chromatography with tris(2,2'-bipyridyl)ruthenium(II)-electrogenerated chemiluminescence detection, *J. Chromatogr. B*, **1995**, *665*, 27-36.
- [58] W. Y. Lee, T. A. Nieman, Evaluation of use of tris(2,2'-bipyridyl)ruthenium(III) as a chemiluminescent reagent for

- quantitation in flowing streams, *Anal. Chem.*, **1995**, *67*, 1789-1796.
- [59] J. P. Preston, T. A. Nieman, An electrogenerated chemiluminescence probe and its application utilizing tris(2,2'-bipyridyl)ruthenium(II) and luminol chemiluminescence without a flowing stream, *Anal. Chem.*, **1996**, *68*, 966-970.
- [60] D. R. Skotty, W. Y. Lee, T. A. Nieman, Determination of dansyl amino acids and oxalate by HPLC with electrogenerated chemiluminescence detection using tris(2,2'-bipyridyl)ruthenium(II) in the mobile phase, *Anal. Chem.*, **1996**, *68*, 1530-1535.
- [61] Z. He, H. Gao, L. Yuan, Q. Luo, Y. Zeng, Simultaneous determination of oxalic and tartaric acid with chemiluminescence detection, *Analyst*, **1997**, *122*, 1343-1346.
- [62] F. Li, H. Gui, X. Q. Lin, Potential-resolved electrochemiluminescence of $\text{Ru}(\text{bpy})_3^{2+}/\text{C}_2\text{O}_4^{2-}$ system on gold electrode, *Luminescence*, **2002**, *17*, 117-122.
- [63] X. Zhang, A. J. Bard, Electrogenerated chemiluminescent emission from an organized (L-B) monolayer of a tris(2,2'-bipyridine)ruthenium(2+)-based surfactant on semiconductor and metal electrodes, *J. Phys. Chem.*, **1988**, *92*, 5566-5569.
- [64] C. J. Miler, P. Mc Cord, A. J. Bard, Study of langmuir monolayers of ruthenium complexes and their aggregation by electrogenerated chemiluminescence, *Langmuir*, **1991**, *7*, 2781-2787.
- [65] Y. S. Obeng, A. J. Bard, Electrogenerated chemiluminescence. 53. Electrochemistry and emission from adsorbed monolayers of a tris(bipyridyl)ruthenium(II)-based surfactant on gold and tin oxide electrodes, *Langmuir*, **1991**, *7*, 195-201.
- [66] Y. Sato, K. Uosaki, Electrochemical and electrogenerated chemiluminescence properties of tris(2,2'-bipyridine)ruthenium(II)-tridecanethiol derivative on ITO and

- gold electrodes, *J. Electroanal. Chem.*, **1995**, 384, 57-66.
- [67] I. Rubinstein, A. J. Bard, Unique properties of chromophore-containing bilayer aggregates: enhanced chirality and photochemically induced morphological change, *J. Am. Chem. Soc.*, **1980**, 102, 6642-6644.
- [68] C. R. Martin, I. Rubinstein, A. J. Bard, Polymer films on electrodes. 9. Electron and mass transfer in Nafion films containing tris(2,2'-bipyridine)ruthenium(2+), *J. Am. Chem. Soc.*, **1982**, 104, 4817-4824.
- [69] O. Dvorak, M. K. De Armond, Electrode modification by the sol-gel method, *J. Phys. Chem.*, **1993**, 97, 2646-2648.
- [70] C. Z. Zhao, N. Egashira, Y. Kurauchi, K. Ohga, Electrochemiluminescence oxalic acid sensor having a platinum electrode coated with chitosan modified with a ruthenium (II) complex, *Electrochim. Acta*, **1998**, 43, 2167-2173.
- [71] A. N. Khramov, M. M. Collinson, Electrogenenerated chemiluminescence of tris(2,2'-bipyridyl)ruthenium(II) ion-exchanged in Nafion-silica composite films, *Anal. Chem.*, **2000**, 72, 2943-2948.
- [72] H. Wang, G. Xu, S. Dong, Electrochemiluminescence of tris(2,2'-bipyridine)ruthenium(II) immobilized in poly(*p*-styrenesulfonate)-silica-Triton X-100 composite thin-films, *Analyst*, **2001**, 126, 1095-1099.
- [73] H. Wang, G. Xu, S. Dong, Electrochemiluminescence of tris(2,2'-bipyridyl)ruthenium (II) ion-exchanged in polyelectrolytesilica composite thin-films, *Electroanalysis*, **2002**, 14, 853-857.
- [74] W. Y. Lee, H. N. Choi, S. H. Cho, Electrogenenerated chemiluminescence from tris(2,2'-bipyridyl)ruthenium(II) immobilized in titania-perfluorosulfonated ionomer composite films, *Anal. Chem.*, **2003**, 75, 4250-4256.
- [75] H. Wang, G. Xu, S. Dong, Electrochemiluminescence sensor using

- tris(2,2'-bipyridyl)ruthenium(II) immobilized in Eastman-AQ55D–silica composite thin-films, *Anal. Chim. Acta*, **2003**, *480*, 285-290.
- [76] W. Cao, J. Jia, X. Yang, S. Dong, E. Wang, Capillary electrophoresis with solid-state electrochemiluminescence detector *Electrophoresis*, **2002**, *23*, 3692-3698.
- [77] M. Bida, F. G. Gao, A. J. Bard, Electrochemistry and electrogenerated chemiluminescence of a thin solid film of a hydrophobic tris(bipyridine) Ru(II) derivative in contact with an aqueous solution, *J. Solid State Electrochem.*, **2004**, *8*, 706-717.
- [78] Y. Du, H. Wei, J. Z. Kang, J. L. Yan, X. B. Yin, X. R. Yang, E. K. Wang, Microchip capillary electrophoresis with solid-state electrochemiluminescence detector, *Anal. Chem.*, **2005**, *77*, 7993-7997.
- [79] J. Li, Y. H. Xu, H. Wei, T. Huo, E. K. Wang, Electrochemiluminescence sensor based on partial sulfonation of polystyrene with carbon nanotubes, *Anal. Chem.*, **2007**, *79*, 5439-5443.
- [80] X. P. Sun, Y. Du, L. X. Zhang, S. J. Dong, E. K. Wang, Luminescent supramolecular microstructures containing Ru(bpy)₃²⁺: solution-based self-assembly preparation and solid-state electrochemiluminescence detection application, *Anal. Chem.*, **2007**, *79*, 2588-2592.
- [81] H. Wang, G. Xu, S. Dong, Electrochemistry and electrochemiluminescence of stable tris(2,2'-bipyridyl)ruthenium(II) monolayer assembled on benzene sulfonic acid modified glassy carbon electrode, *Talanta*, **2001**, *55*, 61-67.
- [82] M. Sykora, T. J. Meyer, Electrogenerated chemiluminescence in SiO₂ sol-gel polymer composites, *Chem. Mater.*, **1999**, *11*, 1186-1189.
- [83] J. M. Clear, J. M. Kelly, D. C. Pepper, J. G. Vos, Preparation, reactions and photoreaction of bis(bipyridyl)poly-4-vinylpyridineruthenium(II) complexes, *Inorg. Chim. Acta*, **1979**, *33*, L139-140.

- [84] O. Haas, J. G. Vos, Preparation, characterisation and electrochemistry of electrodes coated with poly (4-vinylpyridine) bound ruthenium complexes, *J. Electroanal. Chem.*, **1980**, *113*, 139-149.
- [85] J. M. Calvert, T. J. Meyer, Poly(pyridyl)ruthenium(II) complexes of poly(4-vinylpyridine). Synthesis, characterization, and investigation of optical and electrochemical properties, *Inorg. Chem.*, **1981**, *20*, 27-33.
- [86] O. Haas, M. Kriens, J. G. Vos, An electrochemical investigation of photosubstitution processes at glassy-carbon electrodes coated with polymer-bound ruthenium complexes, *J. Am. Chem. Soc.*, **1981**, *103*, 1318-1319.
- [87] J. M. Calvert, J. V. Caspar, R. A. Binstead, T. D. Westmoreland, T. J. Meyer, Metallopolymer photochemistry. Photophysical, photochemical and photoelectrochemical properties of (bpy)₂RuII sites bound to poly(4-vinylpyridine), *J. Am. Chem. Soc.*, **1982**, *104*, 6620-6627.
- [88] A. P. Doherty, M. A. Stanley, D. Leech, J. G. Vos, Oxidative detection of nitrite at an electrocatalytic [Ru(bipy)₂poly-(4-vinylpyridine)₁₀Cl]Cl electrochemical sensor applied for the flow injection determination of nitrate using a Cu/Cd reductor column, *Anal. Chim. Acta*, **1996**, *319*, 111-120.
- [89] R. J. Forster, C. F. Hogan, Mediated electron transfer for electroanalysis: transport and kinetics in tin films of [Ru (bpy)₂PVP₁₀] (ClO₄)₂, *Anal. Chim. Acta*, 1999, *396*, 13-21.
- [90] C. F. Hogan, R. J. Forster, Electrochemiluminescent metallopolymer coatings: combined light and current detection in flow injection analysis, *Anal. Chem.*, **2000**, *72*, 5576-5582.
- [91] A. Mugweru, J. F. Rusling, Catalytic square-wave voltammetric detection of DNA with reversible metallopolymer-coated electrodes, *Electrochem. Comm.*, 2001, *3*, 406-409.
- [92] A. Mugweru, J. F. Rusling, Square wave voltammetric detection of

- chemical DNA damage with catalytic poly(4-Vinylpyridine)-Ru(bpy)₂²⁺ films, *Anal. Chem.*, **2002**, *74*, 4044-4049.
- [93] B. Wang, J. F. Rusling, Voltammetric sensor for chemical toxicity using [Ru(bpy)₂poly(4-vinylpyridine)₁₀Cl]⁺ as catalyst in ultrathin films. DNA damage from methylating agents and an enzyme-generated epoxide, *Anal. Chem.*, **2003**, *75*, 4229-4235.
- [94] L. Dennany, R. J. Forster, J. F. Rusling, Simultaneous direct electrochemiluminescence and catalytic voltammetry detection of DNA in ultrathin films, *J. Am. Chem. Soc.*, 2003, *125*, 5213-5218.
- [95] S Iijima, Helical microtubules of graphitic carbon., *Nature (London)*, **1991**, *351*, 56.
- [96] K. Gong, Y. Yan, M. Ahang, L. Su, S. Xiong, L. Mao, Electrochemistry and electroanalytical applications of carbon nanotube : a review, *Anal. Sci.*, 2005, *21*, 1383-1393.
- [97] A. Merkoci, M. Pumera, X. Liopis, B. Perez, M. del Valle, S. Alegret, New materials for electrochemical sensing VI : Carbon nanotubes, *Trends Anal. Chem.*, **2005**, *24*, 826-838.
- [98] T. W. Ebbesen, P. M. Ajayan, Large-scale synthesis of carbon nanotubes, *Nature (London)*, **1992**, *358*, 220-222.
- [99] Y. Ando, S. Iijima, Preparation of carbon nanotubes by arc-discharge evaporation, *Jpn. J. Appl. Phys.*, **1993**, *32*, 107-109.
- [100] T. Guo, P. Nikolaev, A. Thess, D. T. Colbert, R. E. Smalley, Catalytic growth of single-walled nanotubes by laser vaporization, *Chem. Phys. Lett.*, **1995**, *243*, 49-54.
- [101] P. L. Walker Jr., J. F. Rakszawski, G. R. Imperial, Carbon formation from carbon monoxide-hydrogen mixtures over iron catalysts. I. Properties of carbon formed, *J. Phys. Chem.*, **1959**, *63*, 133-140.
- [102] N. Nakashima, Soluble carbon nanotubes: Fundamentals and

- applications, *International Journal of Nanoscience*, **2005**, *4*, 119-137.
- [103] F. Valentini, A. Amine, S. Orlanducci, M. L. Terranova, G. Palleschi, Carbon nanotube purification: Preparation and characterization of carbon nanotube paste electrodes, *Anal. Chem.*, **2003**, *75*, 5413-5421.
- [104] J. Wang, M. Musameh, Y. Lin, Solubilization of carbon nanotubes by Nafion toward the preparation of amperometric biosensors, *J. Am. Chem. Soc.*, **2003**, *125*, 2408-2409.
- [105] G. A. Rivas, M. D. Rubianes, Carbon nanotubes paste electrode, *Electrochem. Commun.*, **2003**, *5*, 689-694.
- [106] M. Olek, J. Ostrander, S. Jurga, H. Mohwald, N. Kotov, K. Kempa, M. Giersig, Layer-by-layer assembled composites from multiwall carbon nanotubes with different morphologies, *Nano Lett.*, **2004**, *4*, 1889-1895.
- [107] S. Qin, D. Qin, W. T. Ford, J. E. Herrera, D. E. Resasco, Grafting of poly(4-vinylpyridine) to single-walled carbon nanotubes and assembly of multilayer films, *Macromolecules*, **2004**, *37*, 9963-9967.
- [108] J. H. Rouse, P. T. Lillehei, Electrostatic assembly of polymer/single walled carbon nanotube multilayer films, *Nano Lett.*, **2003**, *3*, 59-62.
- [109] A. A. Mamedov, N. A. Kotov, M. Prato, D. M. Guldi, J. P. Wicksted, A. Hirsch, Molecular design of strong single-wall carbon nanotube/polyelectrolyte multilayer composites, *Nat. Mater.*, **2002**, *1*, 190-194.
- [110] M. Zhang, K. Gong, H. Zhang, L. Mao, Layer-by-layer assembled carbon nanotubes for selective determination of dopamine in the presence of ascorbic acid, *Biosensors and Bioelectronics*, **2005**, *20*, 1270-1276.
- [111] C. E. Banks, R. R. Moore, T. J. Davies, R. G. Compton; Investigation of modified basal plane pyrolytic graphite electrodes: definitive evidence for the electrocatalytic properties of the ends of carbon nanotubes, *Chem. Commun.*, **2004**, *16*, 1804-1805.

- [112] J. J. Gooding, R. Wibowo, J. Liu, W. Yang, D. Losic, S. Orbons, F. J. Mearns, J. G. Shapter, D. B. Hibbert, Protein electrochemistry using aligned carbon nanotube arrays, *J. Am. Chem. Soc.*, **2003**, *125*, 9006-9007.
- [113] J. M. Nugent, K. S. V. Santhanam, A. Rudio, P. M. Ajayan, Fast electron transfer kinetics on multiwalled carbon nanotube microbundle electrodes, *Nano Lett.*, **2001**, *1*, 87-91.
- [114] R. L. McCreery, A. J. Bard, *Electroanalytical Chemistry* 2ed., **1991**, Dekker, New York.
- [115] P. J. Britto, K. S. V. Santhanam, P. J. Ajayan, Carbon nanotube electrode for oxidation of dopamine, *Bioelectrochem. Bioenerg.*, 1996, *41*, 121-125.
- [116] H. Luo, Z. Shi., N. Li, Z. Gu, Q. Zhuang, Investigation of the electrochemical and electrocatalytic behavior of single-wall carbon nanotube film on a glassy carbon electrode, *Anal. Chem.*, **2001**, *73*, 915-920.
- [117] R. R. Moore, C. E. Banks, R. G. Compton, Basal plane pyrolytic graphite modified electrodes: Comparison of carbon nanotubes and graphite powder as electrocatalysts, *Anal. Chem.*, **2004**, *76*, 2677-2682.
- [118] C. E. Banks, T. J. Davies, G. G. Wildgoose, R. G. Compton, Electrocatalysis at graphite and carbon nanotube modified electrodes: edge-plane sites and tube ends are the reactive sites, *Chem. Commun.*, **2005**, *7*, 829-841.
- [119] J. J. Gooding, A. Chou, T. Bocking, N. K. Singh, Demonstration of the importance of oxygenated species at the ends of carbon nanotubes for their favourable electrochemical properties, *Chem. Commun.*, **2005**, *1*, 842-844.
- [120] M. Musameh, N. S. Lawrence, J. Wang, Electrochemical activation of carbon nanotubes, *Electrochem. Commun.*, 2005, *7*, 14-18.

- [121] A. W. Bott, Electrochemical techniques for the characterization of redox polymers, *Current Separations*, **2001**, *19*, 71-75
- [122] A. J. Bard, L. R. Faulkner, Electrochemical methods : fundamentals and applications, Wiley, New York, 2001.
- [123] S. Treimer, A. Tang, D. C. Johnson, A consideration of the application of Koutecký-Levich plots in the diagnosis of charge-transfer mechanisms at rotated disk electrodes, *Electroanalysis*, **2002**, *14*, 165-171.
- [124] W. J. Albery, A. R. Hillman, Transport and kinetics in modified electrodes, *J. Electroanal. Chem.*, **1984**, *170*, 27-49.
- [125] 盧玫純，Bis(2,2'-bipyridyl)ruthenium(II)-poly(4-vinylpyridine)修飾ITO電極之電化學行為及電激發化學發光研究，東海大學碩士論文，2003。
- [126] J. M. Clear, J. M. Kelly, D. C. Pepper, J. G. Vos, Preparation, reactions and photoreaction of bis(bipyridyl)poly-4-vinylpyridineruthenium(II) complexes, *Inorg. Chim. Acta*, **1979**, *33*, L139-L140.
- [127] M. D. Rubianes, G. A. Rivas, Carbon nanotubes paste electrode, *Electrochem. Commun.*, **2003**, *5*, 689-694.
- [128] F. Valentini, S. Orlanducci, M. L. Terranova, A. Amine, G. Palleschi, Carbon nanotubes as electrode materials for the assembling of new electrochemical biosensors, *Sens. Actuators, B*, **2004**, *100*, 117-125.
- [129] K. Wu, J. Fei, S. Hu, Simultaneous determination of dopamine and serotonin on a glassy carbon electrode coated with a film of carbon nanotubes, *Anal. Biochem.*, **2003**, *318*, 100-106.
- [130] W. C. Poh, K. P. Loh, W. De Zhang, S. Triparthy, J. Ye, F. Sheu, Biosensing properties of diamond and carbon nanotubes, *Langmuir*, **2004**, *20*, 5484-5492.
- [131] K. Gong, M. Zhang, Y. Yan, L. Su, L. Mao, S. Xiong, Y. Chen, Sol-gel-derived ceramic-carbon nanotube nanocomposite electrodes:

- tunable electrode dimension and potential electrochemical applications, *Anal. Chem.*, **2004**, *76*, 6500-6505.
- [132] J. Wang, M. Musameh, Carbon nanotube/teflon composite electrochemical sensors and biosensors, *Anal. Chem.*, **2003**, *75*, 2075-2079.
- [133] M. Musameh, N. S. Lawrence, J. Wang, Electrochemical activation of carbon nanotubes, *Electrochem. Commun.*, **2005**, *7*, 14-18.
- [134] R. Antiochia, I. Lavagnini, P. Pastore, A comparison between the use of a redox mediator in solution and of surface modified electrodes in the electrocatalytic oxidation of nicotinamide adenine dinucleotide, *Bioelectrochemistry*, **2004**, *64*, 157-163.
- [135] J. Chen, J. Bao, C. Cai, T. Lu, Electrocatalytic oxidation of NADH at an ordered carbon nanotubes modified glassy carbon electrode, *Anal. Chim. Acta*, **2004**, *516*, 29-34.
- [136] M. Musameh, J. Wang, A. Merkoci, Y. Lin, Low-potential stable NADH detection at carbon-nanotube-modified glassy carbon electrodes, *Electrochem. Commun.*, **2002**, *4*, 743-746.
- [137] J. Ye, Y. Wen, W. Zhang, Nonenzymatic glucose detection using multi-walled carbon nanotube electrodes, *Electrochem. Commun.*, **2004**, *6*, 66-70.
- [138] Z. Guo, S. Dong, Electrogenated chemiluminescence from $\text{Ru}(\text{Bpy})_3^{2+}$ ion-exchanged in carbon nanotube/perfluorosulfonated ionomer composite films, *Anal. Chem.*, **2004**, *76*, 2683-2688.
- [139] F. Frehill, M. Panhuis, N. A. Young, W. Henry, J. Hjelm, J. G. Vos, Microscopy and spectroscopy of interactions between metallopolymers and carbon nanotube, *J. Phys. Chem. B*, **2005**, *109*, 13205-13209.
- [140] J. Li, J. D. Qiu, J. J. Xu, H. Y. Chen, X. H. Xia, The synergistic effect of prussian-blue-grafted carbon nanotube/poly(4-vinylpyridine)composites

- for amperometric sensing, *Adv. Funct. Mater.*, **2007**, *17*, 1574-1580.
- [141] L. Zhang, S. Dong, Preparation, reactions and photoreaction of Bis(bipyridyl)poly-4-vinylpyridineruthenium(II) complexes, *Electrochem. Commun.*, **2006**, *8*, 1687-1691.
- [142] M. Zorzi, P. Pastore, F. Magno, A Single Calibration Graph for the Direct Determination of Ascorbic and Dehydroascorbic Acids by Electrogenerated Luminescence Based on $\text{Ru}(\text{bpy})_3^{2+}$ in Aqueous Solution, *Anal. Chem.*, **2000**, *72*, 4934-4939.