

伍、文獻參考

- 莊少鈞。2005。大豆蛋白於能量調節之潛力。食品工業。11:33-54。
- Anderson, J.M., Johnstone, B.M., Kimura, Y. and Kambara, H. (1995 a) Soy protein isolate and its hydrolysate reduce body fat of dietary obese rats and genetically obese mice(yellow KK). *J. Nutr.* 16:349-354.
- Anderson, J.W., Johnstone, E.M. and Cook-Newel, M.E. (1995 b) Meta-analysis of the effects of soy protein intake on serum lipid. *N. Engl. J. Med.* 333:276-282.
- Aoyama, T., Fukui, K., Nakamori, T., Hashimoto, Y., Yamamoto, T. and Takamatsu, K. (2000) Effect of soy and milk whey protein isolates and their hydrolysates on weight reduction in genetically obese mice. *Biosci. Biotechnol. Biochem.* 64: 2594-2600.
- Aoyama, T., Kohono, M., Saito, T., Fukui, K., Takamatsu, K., Yamamoto, T., Hashimoto, Y., Hirtosuka, M. and Kito, M. (2001) Reduction by phytate-reduced soybean β -conglycinin of plasma triglyceride level of young and adult rats. *Biosci. Biotechnol. Biochem.* 65:1071 - 1075.
- Aquila, H., Link, T.A. and Klingenberg M. (1985) The uncoupling protein from brown fat mitochondria is related to the mitochondrial ADP/ATP carrier. Analysis of sequence homologies and of folding of the protein in the membrane. *EMBO J* 4: 2369 - 2376
- Arner, P. (2005) Human fat cell lipolysis: Biochemistry, regulation and clinical role. *Best Pract Res Clin Endocrinol Metab.* 19: 471 - 482.
- Ascencio, C., Torres, N., Isoard-Acosta, F., Gomezperez, F.J., Hernandez-Pando, R. and Tovar, A.R. (2004) Soy protein affects serum insulin and hepatic SREBP-1 mRNA and reduces fatty liver in rats. *J. Nutr.* 134:522-529.
- Barak, Y., Nelson, M. C., Ong, E. S., Jones, Y. Z., Ruiz-Lozano, P., Chien, K. R., Koder, A. and Evans, R. M. *Mol. Cell* (1999) 4:585 - 595.
- Benito, M. and Lorenzo, M. (1992) Hormonal regulation of malic enzyme and glucose-6-phosphate-dehydrogenase expression in fetal brown-adipocyte primary cultures under non-proliferative conditions. *Eur. J. Biochem.* 203: 313 - 319.
- Brasaemle, D. L., Levin, D. M., Adler-Wailes, D.C. and Londos, C. (2000) The lipolytic stimulation of 3T3-L1 adipocytes promotes the translocation of hormone-sensitive lipase to the surfaces of lipid storage droplets. *Biochim. Biophys. Acta.* 1483:251-262.
- Cammalleri, C. and Germinario, R.J. (2003) The effects of protease inhibitors on basal and insulin-stimulated lipid metabolism, insulin

- binding, and signaling. *J. Lipid Res.* 44:103-108.
- Carey G.B. (1998) Mechanisms regulating adipocyte lipolysis. *Adv. Exp. Med. Biol.* 441: 157 - 170.
- Carroll, K. K. (1991) Review of clinical studies on cholesterol-lowering response to soy protein. *J. Am. Diet. Assoc.* 91: 820 - 827.
- Casteilla, L., Champigny, O., Bouillaud, F., Robelin, J. and Ricquier, D. (1989) Sequential changes in the expression of mitochondrial protein mRNA during the development of brown adipose-tissue in bovine and ovine species sudden occurrence of uncoupling protein mRNA during embryogenesis and its disappearance after birth. *Biochem. J.* 257:665-671.
- Chen J., Suetsuna, K. and Yamauchi F. (1995) Isolation and characterization of immunostimulative peptides from soybean. *J Nutr. Biochem.* 6:310 - 313.
- Chernick, S. S., Spooner, P. M., Garrison, M. M. and Scow, R. O. (1986) Effect of epinephrine and other lipolytic agents on intra cellular lipolysis and lipoprotein lipase activity in 3T3-L1 adipocytes. *J. Lipid Res.* 27: 286-294.
- Chiang, W.D., Tsou, M. J., Tsai, Z. Y. and Tsai, T. C. (2006) Angiotensin I-converting enzyme inhibitor derived from soy protein hydrolysate and produced by using membrane reactor. *Food Chem.* 98:725 - 732.
- Choi, I., Park, Y., Choi, H. and Lee, E. H. (2006) Anti-adipogenic activity of rutin in 3T3-L1 cells and mice fed with high-fat diet. *Biofactors.* 26:273-81.
- Christian A. (2005) Drevon. Resistin expression in 3T3-L1 adipocytes is reduced by arachidonic acid. *J. Lipid Res.* 46:143-153.
- Church, F. S., Swaisgood, H. E., Porter, D. H. and Catignani, G. L. (1983) Spectrophotometric assay using o-phthaldialdehyde for determination of proteolysis in milk and isolated milk proteins. *Journal of Dairy Science* 66: 1219-1227.
- Corredig, M. and Dalgleish, D. G. (1997) Studies on the susceptibility of membrane-derived proteins to proteolysis as related to changes in their emulsifying properties. *Food Res. Int.* 30: 689 - 697.
- Danielle, A. J., Kerckhoffs, M., Fred Brouns, Gerard Hornstra and Ronald Mensink P. (2002) Effects on the Human Serum Lipoprotein Profile of α -Glucan, soy protein and isoflavones, plant sterols and stanols, garlic and tocotrienols. *J. Nutr.* 132: 2494 - 2505
- Eleanor, N.W. and Sharon, R.R. (1999) *Understanding Nutrition.* Eight

- edition. An international Thomson Publishing Company. pp 231-289.
- FitzGerald, R.J. and Meisel, H. (2000) Milk protein-derived peptide inhibitors of angiotensin-I-converting enzyme. *J. Nutr.* 84:33- 37.
- Fontaine, K.R. , Yang, D., Gadbury, G.L., Heshka, S., Schwartz, L.G., Murugesan, R., Kraker, J.L., Heo, M., Heymsfield, S.B. and Allison, D.B. (2003) Results of soy-based meal replacement formula on weight, anthropometry, serum lipids & blood pressure during a 40-week clinical weight loss trial. *J. Nutr.* 2:14-20.
- Fortier, M., Soni, K., Laurin, N., Wang, S.P., Mauriège, P., Jirik, F.R., and Mitchell, G.A. (2005) Human hormone-sensitive lipase (HSL): expression in white fat corrects the white adipose phenotype of HSL-deficient mice. *J. Lipid Res.* 46:1860 - 1867.
- Frayn, K.N. and Karpe, F. (2003) Integrative physiology of human adipose tissue. *International Int. J. Obes. Relat. Metab. Disord.* 27: 875 - 888.
- Fredrikson G., Stralfors P., Nilsson N.O. and Belfrage P. (1981) Hormone-sensitive lipase of rat adipose tissue. Purification and some properties. *J. Biol. Chem.* 256:6311-6320.
- Fu, Y., Luo, N., Klein, R.L. and Garvey, W.T. (2005) Adiponectin promotes adipocyte differentiation, insulin sensitivity, and lipid accumulation. *J. Lipid Res.* 46: 1369-1379.
- Fukui, K. , Tachibana, N., Wanezaki, S., Tsuzaki, S., Takamatsu, K., Yamamoto, T., Hashimoto, Y., and Shimoda, T. (2000) Isoflavone-free soy protein prepared by column chromatography reduces plasma cholesterol in rats. *J. Agric. Food Chem.* 50:5717-5721
- Garcia-Jimenez, C., Hernandez, A., Obregon M. J. and Santisteban P. (1993) Malic enzyme gene expression in differentiating brown adipocytes: regulation by insulin and triiodothyronine. *Endocrinology* 132: 1537 - 1543.
- Giorno, L. and Drioli, E. (2000) Biocatalytic membrane reactors: Applications and perspectives. *Trends Biotechnol.* 18:339 - 349.
- Gonzalez-Tello, P., Camacho, F., Jurado, E., Paez, M. P. and Guadex, E. M. (1994) Enzymatic hydrolysis of whey protein II. Molecular -weight range. *Biotechnol. Bioeng.* 44:529 - 534.
- Greaves, K. A. , Parks, J.S. , Williams, J.K. and Wagner, J.D. (1992) Intact dietary soy protein, but not adding an isoflavone -rich soy extract to casein, improves plasma lipids in ovariectomized cynomolgus monkeys. *J. Nutr.* 129: 1585 - 1592.

- Green, H. and Kehinde, O. (1975) An established preadipose cell line and its differentiation in Culture. II. Factors affecting the adipose conversion. *Cell* 5:19-27
- Guadix, A., Camacho, F. and Guadix, E. M. (2006) Production of whey protein hydrolysates with reduced antigenicity in a stable membrane reactor. *J. Food Eng.* 72:398 - 405.
- Halaas, J.L., Gajiwala, K.S., Maffei, M., Cohen, S.L., Chait, B.T., Rabinowitz, D., Lallone, R.L., Burley, S.K. and Friedman J.M. (1995) Weight-reducing effects of the plasma protein encoded by the obese gene. *Science.* 269:543 - 546.
- Haugen, F., Zahid, N., Knut T. Dalen, Hollung, K., Nebb, H. I., Green H. and Kehinde, O. (1975) An established preadipose cell line and its differentiation in culture. II Factors affecting the adipose conversion. *Cell* 5:19 - 27.
- He, J., Jiang, H., Tansey, J.T., Tang, C., Pu, S. and Xu, G. (2006) Calyculin and okadaic acid promote perilipin phosphorylation and increase lipolysis in primary rat adipocytes *Biochim. Biophys. Acta.* 1761:247 - 255.
- Herberg, L., Doppen, W., Major, E. and Gries, F.A. (1974) Dietary-induced hypertrophic- hyperplastic obesity in mice. *J. Lipid Res.* 15:580-585.
- Himms-Hagen J. (1990) Brown adipose tissue thermogenesis: interdisciplinary studies. *FASEB J.* 4:2890-2898.
- Hirsch, J. and Batchelor, B. (1976) Adipose tissue cellularity in human obesity. *Clin. Endocrinol. Metab.* 5:299-311.
- Harmon, A.W. and Harp, J.B. (2001) Differential effects of flavonoids on 3T3-L1 adipogenesis and lipolysis. *Am. J. Physiol. Cell Physiol.* 280:807-813.
- Isley, W.L., Miles, J.M., Patterson, B.W. and Harris, W.S. (2005) The effect of high-dose simvastatin on triglyceride-rich lipoprotein metabolism in patients with type 2 diabetes mellitus. *J. Lipid Res.* 47:193-200.
- Iwai, F. S., Fushiki, T., Tsujikawa, M., Hirose, M., Tsunasawa, S. and Sakiyama, F. (1987) Purification and sequencing of a trypsin-sensitive cholecystokinin-releasing peptide from rat pancreatic juice. Its homology with pancreatic secretory trypsin inhibitor. *J. Biol. Chem.* 262: 8956 - 8959.
- Kambara, H., Motohiko, H., Takamatsu, K. and Kito, M. (2002) Triglyceride lowering effect of soybean β -conglycinin in humans. *Ther. Res.*

23:85 – 89.

- Kammoun, R. , Bejar, S. and Ellouz, R. (2003) Protein size distribution and inhibitory effect of wheat hydrolysates on Neutrased. *Bioresour Technol* 90:249–254.
- Kim, H. S. , Liang, L. , Dean, R. G. , Hausman, D. B. , Hartzell, D. L. and Baile, C. A. (2001) Inhibition of preadipocyte differentiation by myostatin treatment in 3T3-L1 cultures. *Biochem. Biophys. Res. Commun.* 281:902–906.
- Kim, S. Y. , Young, J. J. and Kim S. K. (2001) Purification and characterization of antioxidant peptide from hoki (*Johnius belengerii*) frame protein by gastrointestinal digestion. *J Nutr. Biochem.* 18:31–38.
- Kim, D. W. , Kim, B. S. , Kwon, H. S. , Kim, C. G. , Lee, H. W. , Choi, W. H. and Kim, C. G. (2003) *Exp. Mol. Med.* 5 : 518–526.
- Kim, H. K. , Della-Fera M. , Lin J. and Baile, C. A. (2006) Docosahexaenoic acid inhibits adipocyte differentiation and induces apoptosis in 3T3-L1 preadipocytes. *J. Nutr.* 136:2965–2969.
- Kirk, E. A. , Sutherland, P. , Wang, S. A. , Chait, A. and LeBoeuf, R. C. (1998) Dietary isoflavones reduce plasma cholesterol and atherosclerosis in C57BL/6 mice but not LDL receptor-deficient mice. *J. Nutr.* 128:954–956
- Kong, X. Z. , Zhou, H. and Qian, H. (2006) Enzymatic hydrolysis of wheat gluten by proteases and properties of the resulting hydrolysates. *Food Chem.* 101:615–620.
- Lafontan, M. , Berlan M. and Stich V. (2002) Recent data on the regulation of lipolysis by catecholamines and natriuretic peptides. *Ann. Endocrinol.* 63: 86 – 90.
- Lamas, E. M. , Barros, R. M. , Balcao, V. M. and Malcata, F. X. (2001) Hydrolysis of whey proteins by proteases extracted from *Cynara cardunculus* and immobilized onto highly activated supports. *Enzyme Microb. Technol.* 28:642 – 652.
- Langin, D. , Lucas, S. and Lafontan, M. (2000) Millennium fat-cell lipolysis reveals unsuspected novel tracks. *Horm. Metab. Res.* 32: 443–452.
- Large, V. , Peroni, O. and Letexier, D. (2004) Metabolism of lipids in human white adipocyte. *Diabetes Metab*30: 294 – 309.
- Lasekan, J. B. , Larissa, G. and Sukur, K. (1995) Influence of dietary golden pea proteins versus casein on plasma and hepatic lipids in rats. *Nurs. Res.* 15:71–84.

- Lazar, M. A. (2002) Becoming fat. *Genes Dev.* 16:1 - 5.
- Lichtenstein, A. H. (1998) Soy protein, isoflavones and cardiovascular disease risk. *J. Nutr.* 128: 1589 - 1592.
- Lovati M.R., Manzoni C., Corsini A., Granata A., Frattini R., Fumagalli R. and Sirtori C.R. (1992) Low density lipoprotein receptor activity is modulated by soybean globulins in cell culture. *J. Nutr.* 122:1971 - 1978.
- Mannheim, A. and Cheryan, M. (1990) Continuous hydrolysis of milk protein in a membrane reactor. *J. Food Sci.* 55:381 - 385.
- Manson, J.E., Tosteson, H. and Ridker, P.M. (1992) The primary prevention of myocardial infarction. *N. Engl. J. Med.* 326:1406-1416.
- Manzoni, C., Duranti, M., Eberini, I., Scharnag, H., Marz, W., Castiglioni, S. and Lovati, M.R. (2003) Subcellular localization of soybean 7S globulin in HepG2 cells and LDL receptor up-regulation by its alpha' constituent subunit. *J. Nutr.* 133 : 2149 - 2155.
- Martin-Orue, C., Henry, G. and Bouhallab, S. (1999) Tryptic hydrolysis of j-caseinomacropeptide : Control of the enzymic reaction in a continuous membrane reactor. *Enzyme Microb Technol* 24:173 - 180.
- Maruyama, S., Mitachi, H., Tanaka, H., Tomizuka, N. and Suzuki, H. (1987) Studies on the active site and antihypertensive activity of angiotensin I-converting enzyme inhibitors derived from casein. *Agric. Biol. Chem.* 51:1581 - 1586.
- Meeker, D. R. and Kesten, H. D. (1941) Effect of high protein diets on experimental atherosclerosis of rabbits. *Arch. Pathol.* 31:147-162.
- Mendis, E., Rajapakse, N. and Kim, S.K. (2005) Antioxidant properties of a radicals scavenging peptide purified from enzymatically prepared fish skin gelatin hydrolysate. *J. Agric. Food Chem.* 53:581 - 587.
- Morimoto, C., Tsujita, T. and Okuda, H. (1997) Norepinephrine-induced lipolysis in rat fat cells from visceral and subcutaneous sites: role of hormone-sensitive lipase and lipid droplets. *J. Lipid Res.* 38: 132-138.
- Mueller, E., Drori, S., Aiyer, A., Yie, J., Sarraf, P., Chen, H., Hauser, S., Rosen, E.D., Ge, K., Roeder, R.G. and Spiegelman, B.M. (2002) Genetic analysis of adipogenesis through peroxisome proliferator-activated receptor isoforms. *J. Biol. Chem.* 277: 41925 - 41930.
- Mullally, M.M., Meisel, H. and FitzGerald, R.J. (1997) Angiotensin-I-converting enzyme inhibitory activities of gastric and pancreatic

- proteinasase digests of whey proteins. *International Dairy Journal* 7:299 – 303.
- Nishio, E., Tomiyama, K. , Nakata, H. and Watanabe, Y. (1996) 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor impairs cell differentiation in cultured adipogenic cells(3T3-L1). *Eur. J. Pharmacol.* 301 : 203-206.
- Orlicky, D. J., Lieber, J. G., Morin, C. L. and Evans, R. M. (1998) Synthesis and accumulation of a receptor regulatory protein associated with lipid droplet accumulation in 3T3-L1 cells. *J. Lipid Res.* 39: 1152 – 1161.
- Perea, A. and Ugalde, U. (1996) Continuous hydrolysis of whey proteins in a membrane recycle reactor. *Enzyme Microb. Technol.* 18:29 – 34.
- Pihlanto-Leppälä, A., Rokka, T. and Korhonen, H.. (1998) Angiotensin I converting enzyme inhibitory peptides derived from bovine milk proteins. *International Dairy Journal* 8:325 – 331.
- Pihlanto-Leppälä A. (2000) Bioactive peptides derived from bovine whey proteins: opioid and ACE-inhibitory. *Trends Food Sci. Technol.* 11:347 – 356.
- Potter, S. M. (1995) Overview of proposed mechanisms for the hypocholesterolemic effect of soy. *J. Nutr.* 125: 606 – 611.
- Potter, S. M. (1996) Soy protein and serum lipids. *Curr. Opin. Lipidol.* 7: 260 – 264.
- Potter, S. M. (1998) Soy protein and cardiovascular disease: the impact of bioactive components in soy. *Nutr. Rev.* 56: 231 – 235.
- Prata-Vidal, M., Bouhallab, S., Henry, G. and Aimar, P. (2001) An experimental study of caseinomacropeptide hydrolysis by trypsin in a continuous membrane reactor. *Biochemical Engineering Journal* 8:195 – 202.
- Ren, D., Collingwood, T. N., Rebar, E. J., Wolffe, A. P. and Camp, H. S. (2002) PPAR knockdown by engineered transcription factors: exogenous PPAR2 but not PPAR1 reactivates adipogenesis. *Genes Dev.* 16:27 – 32.
- Ringseis, R., Matthes, B., Lehmann, V., Becker, K., Schöps, R., Ulbrich-Hofmann R. and Eder, K. (2005) Peptides and hydrolysates from casein and soy protein modulate the release of vasoactive substances from human aortic endothelial cells. *Biochim. Biophys. Acta.* 1727:89-97.
- Ross, S. E., Kenneth, N. H., Longo, A., Bennett, C. N., Lucas, P. C., Erickson, R. L. and MacDougald, O. A. (2000) Inhibition of adipogenesis by Wnt

- Signaling. *Science* 289: 950–953.
- Rios, G.M., Belleville, M.P., Paolucci, D. and Sanchez, J. (2004) Progress in enzymatic membrane reactors—A review. *J. Membrane Sci.* 242:189–196.
- Samuelsson, E.G. and Poulsen, O.M. (1987) A peptide preparation, a process for producing it and use of the peptide preparation. European Patent Application A 10:226–221.
- Sato, R., Naguchi, T. and Naito, H. (1986) Casein phosphopeptide (CPP) enhance calcium absorption from the ligated segment of rat small intestine. *J Nutr. Sci. Vitaminol. (Tokyo)* 32:67–76.
- Sen, A., Lea-Currie, Y. R. , Sujkowska, D. , Franklin, D. M., Wilkison, W.O. , Halvorsen, Y. C. and Gimble, J. M. (2001) Adipogenic potential of human adipose derived stromal cells from multiple donors is heterogeneous. *J. Biol. Chem.* 81:312–319.
- Shepherd, P.R., Gnudi, L., Tozzo, E., Yang, H., Leach, F. and Kahn, B.B. (1993) Adipose cell hyperplasia and enhanced glucose disposal in transgenic mice overexpressing GLUT4 selectively in adipose tissue. *J. Biol. Chem.* 268: 22243 – 22246.
- Shimomura, T., Bashmakov, Y., Ikemoto, S., Hotton, J.D., Brown, M.S., and Goldstein, J.L. (1999) Insulin selectively increases SREBP-1c mRNA in the livers of rats with streptozotocin-induced diabetes. *Mol. Cell. Biol.* 19:3760–3768.
- Snyderl, P.B., Esselstyn, J.M., Loughney, K., Wolda, S.L. and Florio, V. (2005) The role of cyclic nucleotide phosphodiesterases in the regulation of adipocyte lipolysis. *J. Lipid Res.* 46:494–503.
- Soppela, P., Nieminen, M., Saarela, S., Keith, J.S., Morrison, J.N., Macfarlane, F. and Trayhurn, P. (1991) Brown fat-specific mitochondrial uncoupling protein in adipose tissues of newborn reindeer. *Am. J. Physiol.* 260: 1229–1234.
- Stralfors, P., Bjorgell, P. and Belfrage, P. (1984) Hormonal regulation of hormone-sensitive lipase in intact adipocytes: Identification of phosphorylated sites and effects on the phosphorylation by lipolytic hormones and insulin, *Proc. Natl. Acad. Sci.* 81:3317–3321.
- Student, A.K., Hsu, R.Y. and Lane, M.D. (1980) Induction of fatty acid synthetase synthesis in differentiating 3T3-L1 preadipocytes. *J. Biol. Chem.* 225:4745–50.
- Suetsuna K., Maekawa K. and Chen J. (2004) Antihypertensive effects of *Undaria pinnatifida* (wakame) peptide on blood pressure in

- spontaneously hypertensive rats. *J. Nutr. Biochem.* 15 :267 - 272.
- Takamatsu, K., Tachibana, N., Matsumoto, I., and Abe, K. (2004) Soy protein functionality and nutrigenomic analysis. *Biofactors* 21:49-53.
- Tardioli, P.W., Sousa, R., Giordano, R. C. and Giordano, R.L.C. (2005) Kinetic model of the hydrolysis of polypeptides catalyzed by Alcalase immobilized on 10% glyoxyl-agarose. *Enzyme Microb. Technol.* 36:555 - 564.
- Tontonoz, P., Hu, E., Graves, R.A., Budavari, A.I. and Spiegelman, B.M. (1994 a) PPAR- γ : tissue-specific regulator of an adipocyte enhancer. *Genes Dev.* 8:1224 - 1234.
- Tontonoz, P., Hu, E. and Spiegelman, B.M. (1994 b) Stimulation of adipogenesis in fibroblasts by PPAR γ 2, a lipid-activated transcription factor. *Cell* 79:1147-1156.
- Trayhurn, P., Thomas, M.E.A. and Keith, J.S. (1993) Postnatal-development of uncoupling protein, uncoupling protein mRNA, and GLUT4 in adipose tissues of goats. *Am. J. Physiol.* 265:676- 682.
- Wang, Y.W., Jones, P.J.H.; Ausman, L.M. and Lichtenstein, A.H. (2004) Soy protein reduces triglyceride levels and triglyceride fatty acid fractional synthesis rate in hypercholesterolemic subjects. *Atherosclerosis* 173:269-275.
- Xu, L., LI, H.M., Huang, Y.B., Wang, H., Nie, G.J. and Zhang, X.Z. (2006) A study on anti-oxidative activity of soybean peptides with linoleic acid peroxidation systems. *Chemical Research in Chinese Universities* 22:205-208.
- Yan, H., Kermouni, A., Abdel-Hafez, M. and Lau, D. C. (2002) Role of cyclooxygenases COX-1 and COX-2 in modulating adipogenesis in 3T3-L1 cells. *J. Lipid Res.* 44:424-429.
- Yoshie, Y. and Wasche, A. (2004) In vitro binding of bile acids by lupin protein isolates and their hydrolysates. *Food Chem.* 88:179 - 184.
- Zhang, Y., Proenca, R., Maffei, M., Barone, M., Leopold, L. and Friedman, J.M. (1994) Positional cloning of mouse obese gene and its human homologue. *Nature* 372:425 - 432.
- Zimmermann R., Strauss J.G. and Haemmerle G. (2004) Fat mobilization in adipose tissue is promoted by adipose triglyceride lipase. *Science* 306: 1383 - 1386.