

第六章 參考文獻

水野 卓、川合正允（賴慶亮譯）(1999) 姑類的化學、生化學，國立編譯館，台北市，第二版。

中國科普博覽，<http://kepu.itsc.cuhk.edu.hk>

王培銘、黃定國（2001）氣動攪拌式發酵槽，食品工業，33(11)：13-26。

王培銘（2002）食藥用菇液態培養製程之開發，食品工業，34(5)：31-35。

李幸儒、陳勁初(1999) 舞茸-溫帶森林中的舞仙子，鄉間小路 25 (4)：32-33。

杜巍、李元瑞、袁靜(2002) 食藥用菌多糖生物活性與鍵結的關係，中國食用菌，21(3)：28-29。

吳經綸、林金枝（1997）灰樹花栽培馴化，食用菌學報 4 (2)：33-39。

吳文騰（2000）發酵槽之設計，生物產業，11(2)：91-97。

吳國暉（2003）以餌料批次發酵進行蟲草菌(CCRC 36421) 多醣生產之影響，大葉大學食品工程研究所，碩士論文。

林家如（2002）浸液發酵培養基與培養條件對藥用茯苓菌絲體及胞外多醣體生成之影響，大葉大學食品工程研究所，碩士論文。

胡琦桂（1994）真菌球狀菌絲體生長之探討，食品工業，26(9)：37-45。

孫希雲、朱明光（1999）灰樹花深層培養基的最優化及一種胞外粗多糖快速測定方法的建立，天津輕工業學院學報，15 (3)：24-28。

陳隆鐘（1997），舞菇生物學之研究(一) 舞菇菌絲生理、生態之研

究，台灣洋菇雜誌。

陳德育、來航線、林雁冰（2004）豬苓的菌絲生長和菌核形成特性及
訓化之研究，食用菌，26(6)：8-9。

郭美娟（1999）菌類之王-舞菇，Nature. Nov-Dec (24) p75-76。

黃仕政、陳勁初（2000）菇類機能性食品之開發，生物產業，
11(3):164-172。

張文菀（2003）不同的液態培養條件對樟芝菌絲生長及多醣形成的
影響，東海大學食品科學研究所，碩士論文。

張志堂（2003）培養條件對靈芝多醣生成及性質之影響，東海大學化
學工程學系研究所，碩士論文。

張毅偉（2003）靈芝中具 - (1 6) 分支之 (1 3) - D - 聚
葡萄糖的性質與檢測，台灣大學食品科技研究所，博士論文。

劉正南、鄭淑芳（1997）中國藥用真菌的現狀和種質資源，中國食用
菌，(6)：30。

劉永銓（1997）生物反應器之放大，化工，44(1)：26-35。

Apar D. K. and Özbek B., (2004) “ α -Amylase inactivation by
temperature during starch hydrolysis.” Process Biochemistry,
39:1137–1144.

Buchalo A. S., Wasser S. p., Reshetnikov S. V. and Grigansky A. P.,
(1999) “Studies on microstructures of vegetative mycelium in the
medicinal mushrooms *Hericium erinaceus* (Bull.:Fr.) pers. And
Grifola frondosa (Dick.:Fr) S. F. Gray (Aphylophoromycetideae).”
Int. J. Med.1 Med. Mushrooms, 1:235-241.

Catley B. J., (1980) “The extracellular polysaccharide,pullulan,
produced by *Aureobasidium pullulans*;a relationship between
elaboration rate and morphology.” J. Gen. Microbiol.,
120:263-268.

Chang T. T., Chou W. N., (2003) "Five polypores (Basidiomycota) new to Taiwan and their cultural characteristics." Bot. Bull. Acad. sin., 44:245-251 .

Chen A. W., Stamets P., Cooper R. B., Huang N. and Han S., (2000) " Ecology, morphology, and morphogenesis in nature of edible and medicinal mushroom *Grifola frondosa* (Dick .:Fr) S. F. Gray -Maitake (AphyllonHoromycetideae)." Int. J. Med. Mushrooms. 2:221-228.

Choi H. S., Cho H. Y., Yang H. C., Ra, K. S. and Suh H. J.,(2001) "Angiotensin I-converting enzyme inhibitor from *Grifola frondosa*." Food Res. Int., 34:177-182.

Costa A., Santoro G., and Asimakopoulos G., (1990) " Cancer chemoprevention - areview of ongoing clinical studies. "Rev. Onc., 3: 657-663.

Dubois M., Gilles K. A., Hamilton J. k., Rebers P. A., Smith F., (1956) "Colorimetric method for determination of sugars and related substances." Anal. Chem., 28:350-356.

FOTKI , <http://public.fotki.com>

Hsu T. H., Shiao L. H., Hsieh C. and Chang D. M., (2002) "A comparison of the chemical composition and bioactive ingredient of the Chinese medicinal mushroom Dong ChongXiaCao, its counterfeit and mimic, and fermented mycelium of *Cordyceps sinensis*." Food chem., 78:463-469.

Janusz M. J., Austen K. F. and Czop J. K. (1989) "Isolation of a yeast heptaglucoside that inhibits mono cyte phagocytosis of zymosan particles [J]." J. Immunol., 142 :959-965.

Jeongseok L., Sang Y. P., Sunwon P. and Anton P. J., (1999) "Control of fed-batch fermentation." Biotechnology Adv., 17:29-48.

Jones K., (1998) "Maitake: A Potent medicinal food." Altern Complement Ther., 4 : 420-429.

Kawagishi H., Nomura A., Mizuno T., Kimura A. and Chiba S., (1990) "Isolation and characterization of a lectin from *Grifola frondosa* fruiting bodies." Biochimica Biophysica Acta., 1034:247-252.

Kawagoe M., Hyakumura K., Suye S., Mili K. and Naoe K., (1997) "Application of bubble column fermentors to submerged culture of *Schizophyllum commune* for production of L-malic acid." J. Fermen. BIOENG., 4:333-336.

Kubo K. and Nanba H., (1997) "Anti-hyperlipidosis effect of Maitake fruit body (*Grifola frondosa*). " J. Biol. Pharm. Bull., 20(7):781-785.

Kulicke W. M., Lettau A. L. and Thielking H., (1997) "Correlation between immunological activity,molar mass, and molecular structure of different (1→3)- β -D-glucans." Carbohydr. Res., 297:135-143.

Lee B. C., (2004) " Submerged culture conditions for the production of mycelial biomass and exopolysaccharides by the edible Basidiomycete *Grifola frondosa*." Enzyme and Microbial Technology, 35:369-376.

Lee B. C., Bae J. T., Pyo H. B., Choe T. B., Kim S. W., Hwang H. J. and Yun J. W. ,(2003) "Biological activities of the polysaccharides produced from submerged culture of the edible Basidiomycete *Grifola frondosa*." Enzyme and Microbial Technology, 32:574-581.

Lee E. W., He P., Kawagishi H. and Sugiyama K., (2000) " Suppression of D-Galactosamine-induced liver injury by mushrooms in rats." Biosci Biotech. Biochem., 64(9):2001-2004.

Lei L. S. and Lin Z. B., (1992) " Effect of Ganoderma polysaccharides on T cell subpopulations and production of interleukin 2 in mixed lymphocyte response." Acta Pharmaceutica Sinica., 27(5): 331-335.

Lieu, C. W., Lee, S. S. and Wang S. Y., (1992) "The effect of *Ganoderma lucidum* on induction of differentiation in leukemic U937 cells." Anticancer Research., 12: 1211-1216.

Litchfield, J. H. , (1979) " Production of single cell protein for usein food and feed. In: Micobial Technology, 2nd ed. " Academic Press, New York. ,93-145.

Maruyama H., Yamazaki K., Murofushi S., Konda C. and Ikekawa T., (1989) "Antitumor activity of *Sarcodon aspratus* (Berk.)S. Ito and *Ganderma licidum*(Fr.)Karst." J. Pharmacobio-Dyn., 12 : 118-123.

Mau J. L., Chang C. N., Huang S. H., Chen C. C., (2004) "Antioxidant properties of methanolic extracts from *Grifola frondosa*,*Morchella esculenta* and *Termitomyces albuminosus* mycelia." Food Chemistry, 87:111-118.

Mau J. L., Lin, H., Ma J. and Song Si-Fu, (2001) "Non-volatile taste components of several speciality mushrooms." Food Chem., 73: 461-466.

Mayell M., (2001) "Maitake extracts and their therapeuticpotential." Altern. Med. Rev., 6(1):48-60.

Miller G.L., (1959) "Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar." Anal. Chen., 31:426.

Mizuno T. , (1995) "Biomolecules of mushroom : Food function and medicinal effect of mushroom fungi." Food Rev. Int., 11:7-21.

Mizuno T., Zhuang C., (1995) "Maitake, *Grifola frondosa* : Pharmacological effects." Food Rev. Int., 11(1):135-149.

Norio Suzuki, Kohzoh Nishibori, Yasuo Oodaira,Shin-ichi Kitamura ,Kenji Michigami ,Keiko Nagata, Yota Tatara ,Byung Rho Lee and Eiji Ichishima , (2005) " Grifolisin, a member of the sedolisin family produced by the fungus *Grifola frondosa*." PHYTOCHEMISTRY, 66:983-990.

Park J. P., Kim S. W., Hwang H. J., Cho Y. J., and Yun J. W., (2002)

"Stimulatory effect of plant oils and fatty acids on the exo-biopolymer production in *Cordyceps militaris*." Enzyme and Microbial technology, 31:250-255.

Roukas T. and Kyriakides M. L., (1999) "Production of pullulan from beet malasses by *Aureobasidium pullulans* in a stirres tank fermentor." J. FoodEng., 40:89-94.

Sasaki T. and Takasuka N., (1976) "Further study of the structure of Lentinan, and anti-tumor polysaccharides from *Lentinus edodes*." Carbohydr. Res., 47:99-104.

Suzuki T., Ohno N., Saito K. and Yadimae T., (1992) "Activation of the complement system by (1→3)- β -D-glucans having different degrees of branching and different ultrastructures." J. Pharmacobio-Dyn., 15:277-285.

Takashi M., (1986) "Fraction and characterization of antitumor polysaccharides from Maritake, *Grifola frondosa*." Agric. Biol. Chem., 50(7):1679-1688.

Tapper H. and Sundler R., (1995) "Glucan receptor and zymosan-induced lysosomal enzyme secretion in macrophages." Biochem. J., 306:829-835.

Tomoda M., Gonda R., Kasahara Y. and Hikino H., (1986) "Autidiabetes drugs. Glycan structure of ganoderans B and D, hypoglycemic glycans of *Ganoderma lucidum* fruit bodies." Phytochem., 25:2817-2820.

Wu J. Z., Cheung Peter C. K., Wong K. H. and Hung N. L., (2003) "Studies on submerged fermentation of *Pleurotus tuber-regium* (Fr.) Singer—Part 1: physical and chemical factors affecting the rate of mycelial growth and bioconversion efficiency." Food Chemistry, 81:389-393.

Wu J. Z., Cheung Peter C. K., Wong K. H. and Hung N. L. (2004) "Studies on submerged fermentation of *Pleurotus tuber-regium* (Fr.) Singer. Part 2: effect of carbon-to-nitrogen ratio of the culture medium on the content and composition of the mycelial dietary fibre." Food Chemistry, 85:101-105.

Yang F. C., Ke Y. F., and Kuo S. S. (2000) "Effect of fatty acids on the mycelial growth and polysaccharide formation by *Ganoderma lucidum* in shake flask cultures." Enzyme and microbial technology, 27:295-301.

Yang Q. Y. and Jong S. C. (1989) "Medicinal mushroom in China." Mush. Sci., 12:631-643.