

參考文獻

- 王伯徹 (1990) 藥用真菌系列報導 (四) 茯苓, 食品工業, 22 (5): 32-38。
- 王利亞、萬惠杰、陳連喜、張紅敏及喻宗源 (1993) 茯苓乙醚萃取物化學成份研究, 中國中藥雜誌, 18 (10): 613-614。
- 王培銘和黃定國 (2001) 氣動攪拌式發酵槽, 食品工業, 33 (11): 13-25。
- 王謙、冀宏、丁方杰、朱朝輝、汪虹、霍紅和顧雅君 (2002) 茯苓的發酵研究, 中國食用菌, 21 (2): 41-42。
- 宋一洋 (2004) 茯苓化學成分及藥理活性之研究 (II), 國防醫學院藥理研究所 碩士論文。
- 宋麟祥、林永浩和沈育維 (2003) 茯苓液態培養技術之開發與應用, 食品工業, 5: 12-21。
- 呂丹妮、余淑娟和曾哲明 (1994) 茯苓對人體血液淋巴球分泌免疫球蛋白的影響, 師大生物學報, 29 (1) 43-51。
- 李得俊、林品汎、陳璿華和王伯徹 (2001) 茯苓液態批次發酵培養最佳化. 中華民國食品科學技術學會會議論文集. 第十六屆第一次.
- 周登成 (1977) 中藥大辭典, 1596-1597。
- 林家如 (2002) 浸液發酵培養基與培養條件對藥用茯苓 (*Wolfiporia cocos*) 菌絲體及胞外多醣體生成之影響, 大葉大學食品工程研究所 碩士論文。
- 吳文騰 (2000) 發酵槽之設計, 生物產業, 11(2): 27-33。
- 吳國暉 (2003) 以饋料批次發酵進行蟲草菌 (CCRC 36421) 多醣生產之影響, 大葉大學食品工程研究所 碩士論文
- 邱紫文和劉懷盛 (1995) 生化反應器概述, 化工技術, 3 (3): 86-92。
- 胡琦桂 (1994) 真菌球狀菌絲體生長之探討, 食品工業, 26(9): 37-45。
- 陳秋桂 (1993) 藥用真菌茯苓調節活性之研究, 國立陽明醫學院藥理學研究所 碩士論文。
- 陳特良 (2002) 饋料批式發酵之進料策略, 化工技術, 10(4): 190-195。
- 黃康泰, (1994) 常用中藥成分與藥裡手冊, 北京, 中國醫藥科技出版社, 1383-1391。
- 黃惠娟 (1996) 茯苓成分對人類肝炎細胞株分泌乙型肝炎表面抗原抑制作用之研究, 國立陽明醫學院藥理學研究所 碩士論文。
- 張志堂 (2003) 培養條件對靈芝多醣生成及性質之影響, 東海大學 碩士論文

- 游英欽 (1996) 以搖瓶振盪及小型發酵槽培養，探討培養基組成及物理化學因子，對多醣生成及生長型態變化的影響，交通大學 生物科技研究所
- 曾美華 (2004) 不同培養條件對舞菇多醣生產之探討，大葉大學 生物產業科技學系
- 楊世海、張紅兵和越英 (1994) 冬蟲夏草研究的新進展，中草藥，25 (9)：488-492。
- 蔡纓和沈忠松 (2001) 茯苓對乙型糖尿病患者的降糖作用，解放軍醫學雜誌，26 (5)：371-372。
- 賴怡琪、劉倩君和曾哲明 (1993) 中藥茯苓對老鼠 B 淋巴球功能的影響，師大生物學報，28：53-63。
- 劉波，1984，中國藥用真菌，山西人民出版社，228。
- 顏正華，中藥學 (上冊)，台北，知音出版社，865-867，1997。
- Ainsworth, G.C., Sparrow, F.K., and Snassman, A.S. (1973) The fungi. Vol. IV A. A Taxonomic Review with Keys: Ascomycetes and Fungi Imperfecti. pp621. Academic Press, Inc. New York.
- Catley, B.J. (1980) The extracellular polysaccharide, pullulan, produced by *Aureobasidium pullulans*; a relationship between elaboration rate and morphology. J. Gen. Microbiol. 120 : 263-268.
- Chen, Y.U., and Chang, H.M., (2004) Antiproliferative and differentiating effect of polysaccharide fraction from fu-ling (*Poria cocos*) on human leukemic U397 and HL-60 cells. Food chem. toxicol. 42, 759-769.
- Cuellar, M. J., Giner, R. M., Recio, M. C., Just, M. Manez, J. S. and Rios, J. L. (1997) Effect of the basidiomycete *Poria cocos* on experimental dermatitis and other inflammatory condition, Chem. Pharm. Bull., 45 (3), 492-494.
- Cukiya, M., Akihisa, T., Hirano, M., Oshikubo, M., Nobukuni, Y., Kimura, Y., Tai, T., Kondo, S., and Nishino, H. (2002) Inhibition of tumor-promoting effect by poricoic acids G and H and other lanostane-type triterpenes and cytotoxic activity of poricoic acids A and G from *Poria cocos*, J. Nat. Prod., 65, 462-465.
- Dobois, M., Gilles, K. A., Hamilton, J. K., Rebers, P.A. and Smith, F. (1956) Colorimetric method for determination of sugars and related substances. Anal Chem. 28(3): 350-356.
- Hattori, T., Hayashi, K., Nagao, T., Furuta, K., Ito, M., and Suzuki, Y. (1992) Antinephritic effect of plant component (3) : Effect of pachyman, a main component of *Poria cocos* Wolf on original-type anti-GBM nephritis in rats and its mechanisms, Jpn. J. Pharmacol., 59, 89-86.
- Hsu, T.H., Shiao, L.H., Hsieh, C., and Chang, D.M. (2002) "A comparison of the chemical composition and bioactive ingredient of the Chinese

- medicinal mushroom Dong ChongXiaCao, its counterfeit and mimic, and fermented mycelium of *Cordyceps sinensis*." Food chem.,78:463-469
- Jeongseok,L., Sang,Y.P., Sunwon,P. and Anton,P.J.(1999) "Control of fed-batch fermentation." Biotechnol.Adv., 17 : 29-48.
- Jin,Y.,Zhang,L.,Chen,Y.,Cheung.P.C.K.,and Chen,L., (2003) Effect of culture media on the chemical and physical characteristics of polysaccharide isolated from *Poria cocos* mycelia. Carbohydr. res.338,1507-1515.
- Kanayama, H., Adachi, N. and Togami, M. (1983) A new antitumor polysaccharide from the mycelia of *Poria cocos* Wolf. Chem. Pharm. Bull. 31(3) : 1115-1118.
- Kanayama,H., Adachi, N., Fukai, Y., Takeuchi, I.,and Togami, M. (1986) Studies on the active polysaccharides from the mycelia of *Poria cocos* Wolf. I. Fractionation and purification of antitumor polysaccharide H11, Yakugaku Zasshi, 106 (3), 199-205.
- Kanematsu ,A.,and Natori,S. (1992) Triterpenoids of Hoelen (fuling), sclerotia of *Poria cocos* Wolf. . 3 β -Hydroxy lanosta-7,9(11),24-Trien-21-oic acid, Chem. Pharm. Bull., 18 (4), 779-783,
- Kaminaga,T.,Yasukawa,K.,and Takido,T. (1996) Inhibitory Effect of *Poria cocos* on 12-*O*-Tetradecanoylphorbol-13-Acetate Induced Ear Oedema and Tumour Promotion in Mouse skin. Phytother. res.,10,581-584
- Kawagoe,M.,Hyakumura,K.,Suye.,Miki,K.,and Naoe,K., (1997) Application of bubble column fermentors to submerged culture of *Schizophyllum commune* for production of L-malic acid.J.Fermen.Bioeng.4 : 333-336.
- Kim,B.S.,Lee,S.C.and Lee,S.Y. (1994) Production of poly by fed-batch culture of *Alcaligenes eu-trophus* with glucose concentration control.Biotech.Bioeng.43 : 892-898.
- Lee ,K. Y., and Jeon ,Y. J. (2003) Polysaccharide isolated from *Poria cocos* sclerotium induces NF-kB/Rel activation and iNOS expression in murine macrophages. International Immunopharmacology. 3, 1353–1362.
- Lee,K. Y., You, H. J., Jeong, H. G., Kang,J. S., Kim, H. M., Rhee,S. D., Jeon, Y. J. (2004) Polysaccharide isolated from *Poria cocos* sclerotium induces NF-nB/Rel activation and iNOS expression through the activation of p38 kinase in murine macrophagesInternational Immunopharmacology 4, 1029–1038.
- Miller G. L. (1959) "Use of ditrosalyscioylic acid reagent for

- determination of reducing sugar.” Anal. Chem., 31: 426.
- McNeil,B.and Harvey,L.M. (1990) Fermentation : A : practical approach.Oxford University Press.
- Nomoto,T.,(1994) Culture of *Poria cocos* Wolf.Japan Patent : JP6277039
- Nukaya, H., Yamashiro,H., Fukazawa, H., Ishida,H ., and Tsuji, K.(1996) Isolation of TPA-induced mouse ear edema from Hoelen, *Poria cocos*, Chem. Pharm. Bull., 44 (4), 847-849.
- Park,J.P., Kim,S.W., Hwang,H.J., Cho,Y.J., and Yun,J.W.(2002)Stimulatory effect of plant oils and fatty acids on the exo-biopolymer production in *Cordyceps militaris*.Enzyme and Microbial Technology, 31 : 250-255.
- Robbins,J.W., and Taylor,K.B.(1989) “Optimization of *Escherichia coli* growth by controlled addition of glucose .” Biotech. Bioeng. 34 : 1289-1294
- Roukass,J.W.,and Kyriakides,M.L. (1998) Production of pullulan from beet malasses by *Aureobasidium pullulans* in a stirred tank fermentor.J.Food Eng.40 : 89-94.
- Simes,J. J., Wootton, H. M., Ralph,B. J., and Pinhey,J. T., (1969) 6 α -Hydroxy-polyporenic acid C, J. Chem. Soc., Chem. Comm., 1150-115.
- Tai,T.,Akahori,A.,and Shingu,T., (1991) Triterpenoid from *Poria cocos*, Phytochemistry, 30 (8), 2796-2797.
- Tai,T.,Akahori,A.and Shingu,T. (1992) Alanosstane triterpenoid from *Poria cocos*,Phytochemistry,31 (7) ,2548-2549,1992.
- Tai,T., Akahori, A., and Shingu,T., (1993) Triterpenoid from *Poria cocos*, Phytochemistry, 32 (5),1239-1244.
- Tai,T. , Akita,Y., Kinoshita, K., Koyama, K., Takahashi,K., and Watanabe,K., (1995a) Triterpenes from the surface lycers of *Poria cocos*, Phytochemistry,39 (5) ,1165-1169.
- Tai.T.,Shingu.T.,Kikochi.T.,Tezuka.Y.,and Akahori A. (1995b)
- Isolation of lanostane-type triterpene acids having an acetoxyl group from sclerotia of *Poria cocos*.Phytochemistry.40 (1) ,225-231.
- Tai.T.,Akita Y.,kinoshita.K.,Takahashi.K.,and Watanabe.K., (1995c) Anti-emetic principles of *Poria cocos* ,Planta Med.,61,527-530,1995.
- Tomohiro,K., Yasukawa,K.,Takaaki,T.,Nunoura,Y.,and Takido.M. (1996) Inhibitory effects of lanostane-type triterpene acids, the components of *Poria cocos*, on tumor promotion by 12- *O* -tetra-decanoylphorbol-13-acetate in two-stage carcinogenesis in

- mouse skin, *Oncology*, 53, 382-385.
- Ukiya, M., Akihisa, T., Hirano, M., Oshikubo, M., Nobukuni, Y., Tai, T., Kondo, S., and Nishino, H., (2002) Inhibition of tumor-promoting effect by poricoic acids G and H and other lanostane-type triterpenes and cytotoxic activity of poricoic acids A and G from *Poria cocos*, *J. Nat. Prod.*, 65, 462-465, 2002.
- Wu, W. T., Huang, T. K., Wang, P. M. and Chen, J. W. (2001) Culture of *Absidia coerulea* for chitinase production in a modified airlift reactor. *J. Chin. Inst. Chem. Engrs.*, 32(3) : 235-240.
- Wu, S. J., Ng, S. J., and Lin, L. T. (2004) Antioxidant Activities of Some Common Ingredients of Traditional Chinese Medicine, *Angelica sinensis*, *Lycium barbarum* and *Poria cocos*. *Phytother. Res.* 18, 1008–1012.
- Yasukawa, K., Kaminaga, T., Kitanaka, S., Tai, T., Nunoura, Y., Natori, S., Takido, S. and M. (1998) 3 β -Hydroxybenzoyldehydrotumulosic acid from *Poria cocos*, and its anti-inflammatory effect, *Phytochemistry*, 48 (8), 1357-1360.
- Yang, F. C. and Hwang, S. Y., (1989) Nutritional studies on submerged culture of *Ganoderma lucidum*. *Tunghai Journal*. 39, 1-10.
- Yang, F. C., Ke, Y. F., and Kuo, S. S. (2000) "Effect of fatty acids on the mycelial growth and polysaccharide formation by *Ganoderma lucidum* in shake flask cultures. *Enzyme and microbial technology*, 27, 295-301
- Zhang, L., Ding, Q., Zhang, P., Zhu, R., and Zhang, Y. (1997) Molecular weight and aggregation behaviour in solution of β -D-glucan from *Poria cocos* sclerotium. *Carbohydr. Res.* 303, 193-197
- Zhang, L., Ding, Q., Diya, M., Ren, L., and Yang, G., Liu, Y. (1999) Investigation of molecular masses and aggregation of β -D-glucan from *Poria cocos* sclerotium by size-exclusion chromatography. *Journal of Chromatography A*, 839, 49-55.