

目 錄

| | |
|---|----|
| 第一章 前言..... | 1 |
| 1.1 研究緣起..... | 1 |
| 1.2 研究目的..... | 2 |
| 第二章 文獻回顧..... | 4 |
| 2.1 纖維廢棄物..... | 4 |
| 2.1.1 來源..... | 4 |
| 2.1.2 處理方法..... | 4 |
| 2.2 纖維素之生物轉化..... | 5 |
| 2.3 嗜熱厭氧纖維分解菌..... | 7 |
| 2.4 甲烷菌..... | 10 |
| 2.4.1 嗜熱性甲烷菌(Thermophilic methanogens)..... | 12 |
| 2.5 <i>Methanobacteriaceae</i> Family 甲烷桿菌科..... | 15 |
| 2.6 <i>Methanothermobacter</i> 屬甲烷菌..... | 17 |
| 2.6.1 <i>Methanothermobacter thermautotrophicus</i> | 17 |
| 2.6.2 <i>Methanothermobacter defluvii</i> | 20 |
| 2.6.3 <i>Methanothermobacter marburgensis</i> | 20 |
| 2.6.4 <i>Methanothermobacter thermoflexus</i> | 20 |
| 2.6.5 <i>Methanothermobacter thermophilus</i> | 21 |
| 2.6.6 <i>Methanothermobacter wolfeii</i> | 21 |
| 2.7 微生物多樣性探討..... | 22 |

| | |
|--|----|
| 2.7.1 微生物分類法..... | 22 |
| 2.7.2 分子分類學..... | 22 |
| 2.7.3 分子分類方法..... | 23 |
| 2.7.3.1 聚合酶連鎖反應(Polymerase Chain Reaction, PCR)..... | 25 |
| 2.7.3.2 變性梯度凝膠電泳(Denaturing Gradient Gel Electrophoresis, DGGE)..... | 29 |
| 第三章 實驗材料與方法..... | 29 |
| 3.1 實驗流程設計..... | 29 |
| 3.2 混合菌種來源..... | 26 |
| 3.3 菌種培養..... | 31 |
| 3.3.1 嗜熱厭氧培養基..... | 31 |
| 3.3.2 Hungate 除氧系統..... | 35 |
| 3.3.3 纖維降解菌群培養..... | 35 |
| 3.4 嗜熱性纖維素降解菌群分析..... | 37 |
| 3.4.1 DNA 萃取方法..... | 37 |
| 3.4.2 萃取之 DNA 產物檢視..... | 38 |
| 3.4.3 DNA 的純度及定量檢測..... | 38 |
| 3.4.4 聚合酶連鎖反應(Polymerase Chain Reaction, PCR)..... | 39 |
| 3.4.5 變性梯度凝膠電泳(Denaturing Gradient Gel Electrophoresis, DGGE)分析菌群結構..... | 43 |

| | |
|---|----|
| 3.4.6 變性梯度凝膠膠片萃取..... | 45 |
| 3.4.7 核酸序列分析..... | 46 |
| 3.5 嗜熱性甲烷菌之純菌分離..... | 47 |
| 3.5.1 無機鹽培養基..... | 48 |
| 3.5.2 嗜熱甲烷菌之 16S rDNA 分析..... | 52 |
| 3.5.2.1 甲烷菌之 DNA 萃取..... | 52 |
| 3.5.2.2 嗜熱甲烷菌之 16S rDNA 擴增與親緣分析..... | 53 |
| 3.6 革蘭氏染色..... | 53 |
| 3.7 菌株型態觀察..... | 55 |
| 3.7.1 位相差顯微鏡(Phase contract microscope)..... | 55 |
| 3.7.2 穿透式電子顯微鏡(Transmission electron microscope, TEM)..... | 55 |
| 3.7.3 菌種保存..... | 56 |
| 3.8 基質利用測試..... | 56 |
| 3.9 生長因子測試..... | 59 |
| 3.10 溫度..... | 59 |
| 3.11 pH 值..... | 59 |
| 3.12 氯化鈉濃度..... | 60 |
| 3.13 抗生素感受測試..... | 60 |

| | |
|---|-----|
| 3.14 SDS 感受性測試..... | 62 |
| 3.15 G+C content (guanine plus cytosine content)分析..... | 62 |
| 3.16 分析方法..... | 64 |
| 3.16.1 生長速率..... | 64 |
| 3.16.2 甲烷濃度分析..... | 64 |
| 第四章 結果與討論..... | 66 |
| 4.1 纖維分解產甲烷混合菌群樣品來源..... | 66 |
| 4.2 核酸萃取所得 DNA 濃度及純度..... | 66 |
| 4.3 PCR 擴增 16S rDNA 片段..... | 68 |
| 4.4 DGGE 進行纖維分解產甲烷混合菌群結構分析..... | 72 |
| 4.4.1 Nested PCR..... | 72 |
| 4.4.2 變性梯度凝膠電泳(DGGE)分析..... | 75 |
| 4.4.3 纖維分解產甲烷混合菌群結構分析..... | 78 |
| 4.5 嗜熱性甲烷菌之純化分離..... | 86 |
| 4.6 分離甲烷菌之定性與鑑定..... | 90 |
| 4.6.1 形態研究..... | 90 |
| 4.6.2 16S rDNA 序列分析..... | 93 |
| 4.6.3 菌株生化特性之探討..... | 97 |
| 4.6.3.1 基質利用..... | 97 |
| 4.6.3.2 生長因子..... | 101 |

| | |
|---|-----|
| 4.6.3.3 基質共利用(co- utilized)..... | 102 |
| 4.6.3.4 最佳生長溫度..... | 105 |
| 4.6.3.5 pH 值..... | 107 |
| 4.6.3.6 NaCl 影響..... | 109 |
| 4.6.3.7 抗生素感受性..... | 111 |
| 4.6.3.8 SDS 感受性..... | 114 |
| 4.6.4 G+C content (guanine plus cytosine content)分析..... | 115 |
| 4.6.5 研究菌株與 <i>Methanothermobacter</i> sp.比較..... | 117 |
| 第五章 結論與建議..... | 120 |
| 5.1 結論..... | 120 |
| 5.2 建議..... | 122 |
| 第六章 參考文獻..... | 124 |
| 附錄..... | 132 |
| 附錄一 Partial sequences of eubacterial 16S rDNA fragments from DGGE gel..... | 132 |
| 附錄二 Partial sequences of archael 16S rDNA fragments from DGGE gel..... | 137 |
| 附錄三 Partial sequence of 16S rDNA of strain THUT3..... | 142 |

表目錄

| | |
|--|----|
| Table 2.1 List of thermophilic anaerobic cellulolytic bacteria..... | 9 |
| Table 2.2 Reaction and standard changes in free energies for methanogenesis..... | 11 |
| Table 2.3 Classification of microorganism by different optimal growth temperatures..... | 13 |
| Table 2.4 List of thermophilic methanogens..... | 14 |
| Table 2.5 Some characteristics of methanogenic archaea of Family <i>Methanobacteriaceae</i> | 18 |
| Table 3.1 Components of TA medium..... | 32 |
| Table 3.2 Components of trace element solution of TA medium..... | 33 |
| Table 3.3 Components of vitamin solution of TA medium..... | 34 |
| Table 3.4 PCR primers for amplification of 16S rDNAs..... | 40 |
| Table 3.5 The reaction mixtures of PCR..... | 41 |
| Table 3.6 The reaction mixtures of nested PCR..... | 42 |
| Table 3.7 Reagent and concentration of DGGE. Concentration range of denaturant..... | 44 |
| Table 3.8 Components of MM medium..... | 49 |
| Table 3.9 Components of trace element solution of MM medium..... | 50 |
| Table 3.10 Components of vitamin solution of MM medium..... | 51 |
| Table 3.11 Procedures and results of each step of gram stain..... | 54 |

| | |
|--|-----|
| Table 3.12 Substrates used for utilization test..... | 57 |
| Table 3.13 Co-utilized substrates used for growth of the isolated methanogen..... | 58 |
| Table 3.14 Characteristics of antibiotics tested in this study and their modes of action..... | 61 |
| Table 4.1 The DNA concentration and purity extracted from the mixed culture in different incubation time..... | 67 |
| Table 4.2 Conditions for PCR amplification of 16S rDNA from cellulose-degrading mixed culture..... | 70 |
| Table 4.3 Conditions for amplifying variable regions of 16S rDNA by nested PCR..... | 73 |
| Table 4.4 Conditions for denature gradient gel electrophoresis of Bacteria and Archaea..... | 76 |
| Table 4.5 The similarity of variable region in 16S rDNA among isolated Bacterial OTU..... | 79 |
| Table 4.6 The similarity of variable region in 16S rDNA among isolated Archaeal OTU..... | 80 |
| Table 4.7 Similarity of 16S rDNA sequence of strain THUT3 to <i>Methanothermobacter</i> sp..... | 94 |
| Table 4.8 Antibiotic resistance of the strain THUT3 to different antibiotics..... | 113 |
| Table 4.9 Susceptibility to SDS lysis..... | 116 |

Table 4.10 Differentiation of strain THUT3 and other
Methanothermobacter species.....119

圖 目 錄

| | |
|--|----|
| Fig. 2.1 Diagram of anaerobic biodegradation of cellulose to methane by microbial communities in soils and freshwater sediments..... | 6 |
| Fig. 2.2 Structure of the universal (U1-U8)and variable region in the SSU rRNAs..... | 24 |
| Fig. 2.3 Schematic diagram of denaturing gradient gel electrophoresis | 28 |
| Fig. 3.1 Flow chart of experimental design in this study..... | 30 |
| Fig. 3.2 Schematic diagram of Hungate gas station..... | 36 |
| Fig. 4.1. Electrophoresis of PCR-amplified bacterial 16S rDNA from cellulose-degrading thermophilic mixed culture..... | 71 |
| Fig. 4.2 Electrophoresis of nested PCR amplified variable region fragment of 16S rDNA..... | 74 |
| Fig. 4.3 DGGE profiles of the thermophilic cellulose-degrading bacterial community and archael community..... | 77 |
| Fig. 4.4 Phylogenic analysis of eubacteria in the thermophilic, anaerobic, cellulose-degrading consortia..... | 82 |
| Fig. 4.5 Phylogenic analysis of archaea in the thermophilic, anaerobic, cellulose-degrading consortia..... | 84 |
| Fig. 4.6. Electrophoresis of PCR-amplified 16S rDNA and variable region fragment..... | 88 |
| Fig. 4.7 Phase-contrast photomicrograph of strain THUT3..... | 91 |
| Fig. 4.8 Transmission electron micrograph of a thin section of whole sell of strain THUT3..... | 92 |

| | |
|---|-----|
| Fig. 4.9 Phylogenic tree of strain THUT3 constructed from 16S rRNA gene sequence. Sequence accession numbers are show in parentheses..... | 96 |
| Fig. 4.10 Growth and methane production rates of strain THUT3 with different substrates in MM medium without yeast extract..... | 99 |
| Fig. 4.11 Growth and methane production rates of strain THUT3 with different substrates in MM medium with yeast extract..... | 100 |
| Fig. 4.12 Growth and methane production strain THUT3 in presence of growth factor (yeast extract or peptone)..... | 103 |
| Fig. 4.13 Growth and methane production rates of strain THUT3 at co-utilized substrates..... | 105 |
| Fig. 4.14 Growth and methane production rates of strain THUT3 at different temperatures..... | 106 |
| Fig. 4.15 Growth and methane production rates of strain THUT3 at different pH..... | 108 |
| Fig. 4.16 Growth and methane production rates of strain THUT3 at different NaCl concentration..... | 110 |